
**A geometrical version of Hardy's
inequality for $\mathring{W}^{1,p}(\Omega)$**

Jesper Tidblom

RESEARCH REPORTS IN MATHEMATICS
NUMBER 16, 2001

DEPARTMENT OF MATHEMATICS
STOCKHOLM UNIVERSITY

Electronic versions of this document are available at
<http://www.matematik.su.se/reports/2001/16>

Date of publication: November 27, 2001
2000 Mathematics Subject Classification: Primary 35P99, Secondary 35P20, 47A75,
47B25.

Postal address:
Department of Mathematics
Stockholm University
S-106 91 Stockholm
Sweden

Electronic addresses:
<http://www.matematik.su.se>
info@matematik.su.se

A geometrical version of Hardy's inequality for $\mathring{W}^{1,p}(\Omega)$

Jesper Tidblom

Abstract

The aim of this article is to prove a Hardy type inequality, concerning functions in $\mathring{W}^{1,p}(\Omega)$ for some domain $\Omega \subset \mathbb{R}^n$, involving the volume of Ω and the distance to the boundary of Ω . The inequality is a generalization of a recently proved inequality by M.Hoffmann-Ostenhof, T.Hoffmann-Ostenhof and A.Laptev [9], which dealt with the special case $p = 2$.

0 Introduction

The history of Hardy type inequalities goes back to Hardy and the 1920's when the following original one-dimensional inequality appeared in [8].

$$\int_0^\infty \left(\frac{F(x)}{x} \right)^p dx \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty f(x)^p dx,$$

where

$$p > 1, \quad f(x) \geq 0 \text{ and } F(x) = \int_0^x f(t) dt$$

(see also [7]). A multidimensional version of this inequality is

$$\int_{\mathbb{R}^n} |\nabla u|^p dx \geq \left| \frac{n-p}{p} \right|^p \int_{\mathbb{R}^n} \frac{|u(x)|^p}{|x|^p} dx, \quad u \in C_0^\infty(\mathbb{R}^n \setminus \{0\}),$$

where $p > 1$ and the constant $\left| \frac{n-p}{p} \right|^p$ is optimal (see for example [13]).

Later on, these inequalities have been generalized and modified in many different ways and the literature concerning such inequalities is extensive. There is an entire book by B.Opic and A.Kufner devoted to various Hardy type inequalities (see [13]). Many other Hardy-Sobolev type inequalities may be found in the excellent book "Sobolev Spaces" [12] by V.G.Maz'ja. In the past few years a lot of articles on the subject has been published, see [3] for a review of recent results in the field. In the article [1] G.Barbatis, S.Filippas and A.Tertikas give a very comprehensive treatment of improved L^p

⁰1991 Mathematics Subject Classification. Primary 35P99; Secondary 35P20, 47A75, 47B25.

Hardy inequalities with best constants, involving various kinds of distance functions.

Let Ω be an open set in \mathbb{R}^n and $\delta(x) = \text{dist}(x, \partial\Omega)$. It is known (see for example [11]) that for any $p > 1$ we have

$$\int_{\Omega} |\nabla u|^p dx \geq c_p \int_{\Omega} \frac{|u|^p}{\delta(x)^p} dx, \quad u \in \mathring{W}^{1,p}(\Omega), \quad (0.1)$$

where Ω is convex and $c_p = \left(\frac{p-1}{p}\right)^p$ is the best constant (see for example [11]). $\mathring{W}^{1,p}(\Omega)$ as usual is the completion of $C_0^\infty(\Omega)$ with respect to the Sobolev norm

$$\|u\|_{W^{1,p}(\Omega)} = \|u\|_{L^p(\Omega)} + \sum_{i=1}^n \left\| \frac{\partial u}{\partial x_i} \right\|_{L^p(\Omega)}.$$

The main result to be proved in this paper is that an extra term of the form

$$\frac{a(p, n)}{|\Omega|^{\frac{p}{n}}} \int_{\Omega} |u(x)|^p dx, \quad (a(p, n) > 0),$$

where $|\Omega| = \text{vol}(\Omega)$, may be added to the right hand side of the inequality (0.1).

Let $\Omega \subset \mathbb{R}^n$ be an arbitrary convex domain. In [2] H.Brezis and M.Marcus proved that the largest possible constant $\lambda(\Omega)$ in the inequality

$$\int_{\Omega} |\nabla u(x)|^2 dx \geq \frac{1}{4} \int_{\Omega} \frac{|u(x)|^2}{\delta^2(x)} dx + \lambda(\Omega) \int_{\Omega} |u(x)|^2 dx, \quad u \in \mathring{W}^{1,2}(\Omega),$$

satisfies

$$\lambda(\Omega) \geq \frac{1}{4 \cdot \text{diam}^2(\Omega)}.$$

In the same paper H.Brezis and M.Marcus have asked whether the above estimate can be replaced by some other estimate of the type $\lambda(\Omega) \geq \alpha |\Omega|^{-2/n}$ for some universal constant $\alpha > 0$.

This question was recently answered affirmative by M.Hoffmann–Ostenhof, T.Hoffmann–Ostenhof and A.Laptev in [9]. In that paper, the following Hardy type inequality

$$\int_{\Omega} |\nabla u(x)|^2 dx \geq \frac{1}{4} \int_{\Omega} \frac{|u(x)|^2}{\delta^2(x)} dx + \frac{\mu_n}{|\Omega|^{\frac{2}{n}}} \int_{\Omega} |u(x)|^2 dx, \quad u \in \mathring{W}^{1,2}(\Omega), \quad (0.2)$$

where

$$\mu_n = \frac{n^{(n-2)/n} |\mathbb{S}^{n-1}|^{2/n}}{4},$$

is established.

Here we shall prove a similar “geometric” inequality for functions from the Sobolev space $\mathring{W}^{1,p}(\Omega)$. More precisely, we shall prove

$$\int_{\Omega} |\nabla u|^p dx \geq c_p \int_{\Omega} \frac{|u(x)|^p}{\delta^p(x)} dx + \frac{a(p, n)}{|\Omega|^{\frac{p}{n}}} \int_{\Omega} |u(x)|^p dx, \quad u \in \mathring{W}^{1,p}(\Omega),$$

where

$$a(p, n) = \frac{(p-1)^{p+1}}{p^p} \cdot \left(\frac{|\mathbb{S}^{n-1}|}{n} \right)^{\frac{p}{n}} \cdot \frac{\sqrt{\pi} \cdot \Gamma\left(\frac{n+p}{2}\right)}{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{n}{2}\right)}.$$

The latter inequality is a generalization of inequality (0.2) for any $p > 1$. In particular, $a(2, n) = \mu_n$.

In section one, we shall, following a method from [9], prove a one-dimensional version of the inequality and in section two, we shall extend it to higher dimensions.

Acknowledgements

I wish to thank my supervisor Ari Laptev for his help and for suggesting the problem.

1 One-dimensional inequalities

Let f be a function defined and differentiable on $(0, b]$ for some $b > 0$. We say that f belongs to the class $\Phi_p(0, b)$ if f is real-valued and there exists a constant $C = C(f)$ such that

$$\sup_{0 < t \leq b} (t^{p-1}|f(t)| + t^p|f'(t)|) \leq C.$$

Throughout this article it is assumed that $p > 1$.

Lemma 1.1. *Let $u \in C^1[0, b]$, $b > 0$, $u(0) = 0$ and let $f \in \Phi_p(0, b)$. Then we have the following inequality :*

$$\int_0^b |u'(t)|^p dt \geq \frac{1}{p^p} \frac{\left| \int_0^b f'(t)|u(t)|^p dt \right|^p}{\left(\int_0^b |f(t) - f(b)|^{\frac{p}{p-1}} |u(t)|^p dt \right)^{p-1}}.$$

Proof. Let c be a constant. We have :

$$\begin{aligned} |(f(b) - c)|u(b)|^p &= \left| \int_0^b f'(t)|u(t)|^p dt \right| \\ &= \left| \int_0^b (f(t) - c)(|u(t)|^p)' dt \right| \\ &= \frac{p}{2} \left| \int_0^b (f(t) - c)(u^{\frac{p}{2}-1} \bar{u}^{\frac{p}{2}} u' + \bar{u}^{\frac{p}{2}-1} u^{\frac{p}{2}} \bar{u}') dt \right| \\ &\leq p \int_0^b |f(t) - c| |u|^{p-1} |u'| dt \\ &\leq p \left(\int_0^b |f(t) - c|^{\frac{p}{p-1}} |u|^p dt \right)^{p-1} \int_0^b |u'|^p dt. \end{aligned}$$

Now put $c = f(b)$ and rise both sides to the power p . We get

$$\left| \int_0^b f'(t)|u|^p dt \right|^p \leq p^p \left(\int_0^b |f(t) - f(b)|^{\frac{p}{p-1}} |u|^p dt \right)^{p-1} \int_0^b |u'|^p dt$$

and we are done. □

Corollary 1.1. *Let u be as in the lemma above and put $f(t) = \frac{t^{1-p}}{1-p} \in \Phi_p(0, b)$. Then the following improved Hardy inequality holds*

$$\begin{aligned} \int_0^b |u'(t)|^p dt &\geq c_p \frac{\left(\int_0^b \frac{|u|^p}{t^p} dt\right)^p}{\left(\int_0^b |t^{1-p} - b^{1-p}|^{\frac{p}{p-1}} |u|^p dt\right)^{p-1}} \\ &\geq c_p \int_0^b \frac{|u|^p}{t^p} dt. \end{aligned}$$

Proof. Use the previous lemma. \square

Now we give a linearized version of the corollary :

Corollary 1.2 (linearized version). *Let u be as above. Then*

$$\int_0^b |u'(t)|^p dt \geq c_p \left(\int_0^b \left(\frac{p}{t^p} - (p-1)(t^{1-p} - b^{1-p})^{\frac{p}{p-1}} \right) |u|^p dt \right). \quad (1.1)$$

Proof. Young's inequality gives us

$$\frac{A^p}{B^{p-1}} \geq pA - (p-1)B.$$

If we put $A = \int_0^b \frac{|u|^p}{t^p} dt$, $B = \int_0^b |t^{1-p} - b^{1-p}|^{\frac{p}{p-1}} |u|^p dt$ and use Corollary 1.1, we get (1.1). \square

An easy consequence of Corollary 1.2 (see also [9]) is

Lemma 1.2. *Let $u \in \mathring{W}^{1,p}(0, 2b)$, $b > 0$. Then we have*

$$\int_0^{2b} |u'(t)|^p dt \geq c_p \int_0^{2b} \left(\frac{p}{\rho(t)^p} - (p-1) \left(\frac{1}{\rho(t)^{p-1}} - \frac{1}{b^{p-1}} \right)^{\frac{p}{p-1}} \right) |u(t)|^p dt,$$

where

$$\rho(t) = \text{dist}(t, \mathbb{R} \setminus [0, 2b]) = \min(t, 2b - t).$$

Proof. By rewriting the inequality (1.1) for the interval $[b, 2b]$ for functions $u \in C^1[b, 2b]$ such that $u(2b) = 0$, we get

$$\int_b^{2b} |u'(t)|^p dt \geq c_p \int_b^{2b} \left(\frac{p}{(2b-t)^p} - \frac{(p-1)}{((2b-t)^{1-p} - b^{1-p})^{\frac{p}{p-1}}} \right) |u|^p dt. \quad (1.2)$$

If we add the inequalities (1.1) and (1.2) and use standard density arguments, we get the statement of the lemma. \square

Theorem 1.1 (one-dimensional version). *Let $u \in \mathring{W}^{1,p}(a, b)$. Then we have*

$$\int_a^b |u'(t)|^p dt \geq c_p \left(\int_a^b \frac{|u(t)|^p}{\rho(t)^p} dt + \frac{p-1}{\left(\frac{b-a}{2}\right)^p} \int_a^b |u(t)|^p dt \right). \quad (1.3)$$

Proof. Without loss of generality we can assume the interval of integration is $[0, 2b]$. The right hand side in Lemma 1.2 may be written

$$c_p \left(\int_0^{2b} \frac{|u(t)|^p}{\rho(t)^p} dt + \int_0^{2b} \frac{p-1}{\rho(t)^p} \left(1 - \left(1 - \left(\frac{\rho(t)}{b} \right)^{p-1} \right)^{\frac{p}{p-1}} \right) |u(t)|^p dt \right).$$

We will now estimate the expression in front of $|u|^p$ in the last integral from below. We begin by noticing that $\rho(t) \leq b$. We get :

$$\begin{aligned} \frac{1}{\rho(t)^p} \left(1 - \left(1 - \left(\frac{\rho(t)}{b} \right)^{p-1} \right)^{\frac{p}{p-1}} \right) &\geq \frac{1}{\rho(t)^p} \left(1 - \left(1 - \left(\frac{\rho(t)}{b} \right)^{p-1} \right) \right) \\ &= \frac{1}{\rho(t)b^{p-1}} \geq \frac{1}{b^p}. \end{aligned}$$

This, together with Lemma 1.2, immediately gives us inequality (1.3). \square

2 Inequalities in higher dimensions

In this section we will extend the one-dimensional results in the previous section to higher dimensions, using almost the same arguments as in [9]. For simplicity I use the same notation as in the mentioned article. If $\nu \in \mathbb{S}^{n-1}$, we put

$$\tau_\nu(x) = \min\{s > 0 : x + s\nu \notin \Omega\}, \quad \rho_\nu(x) = \min(\tau_\nu(x), \tau_{-\nu}(x))$$

$$D_\nu(x) = \tau_\nu(x) + \tau_{-\nu}(x), \quad \Omega_x = \{y \in \Omega : x + t(y-x) \in \Omega, \forall t \in [0, 1]\}$$

$$\delta(x) = \inf_{\nu \in \mathbb{S}^{n-1}} \tau_\nu(x) = \text{dist}(x, \partial\Omega)$$

$d\omega(\nu)$ denotes the normalized surface measure on \mathbb{S}^{n-1} , $\int_{\mathbb{S}^{n-1}} d\omega(\nu) = 1$. Before stating our main theorem we need an auxiliary lemma.

Lemma 2.1.

$$\int_{\mathbb{S}^{n-1}} \left(\frac{2}{D_\nu(x)} \right)^p d\omega(\nu) \geq \left(\frac{n|\Omega_x|}{|\mathbb{S}^{n-1}|} \right)^{-\frac{p}{n}}. \quad (2.1)$$

Proof. Since the function $f(t) = t^{-p}$ is convex when $t > 0$, we can use Jensens inequality to get

$$\int_{\mathbb{S}^{n-1}} \left(\frac{2}{D_\nu(x)} \right)^p d\omega(\nu) \geq \left(\int_{\mathbb{S}^{n-1}} \left(\frac{D_\nu(x)}{2} \right) d\omega(\nu) \right)^{-p}.$$

Since

$$\begin{aligned}
\int_{S^{n-1}} \left(\frac{D_\nu(x)}{2} \right) d\omega(\nu) &= \frac{1}{2} \int_{S^{n-1}} \tau_\nu + \tau_{-\nu} d\omega(\nu) \\
&= \int_{S^{n-1}} \tau_\nu d\omega(\nu) \\
&\leq \left(\int_{S^{n-1}} \tau_\nu^n d\omega(\nu) \right)^{\frac{1}{n}} \\
&= \left(\frac{n|\Omega_x|}{|\mathbb{S}^{n-1}|} \right)^{\frac{1}{n}}
\end{aligned}$$

we obtain (2.1). □

We are now ready for the main theorem.

Theorem 2.1. *Let Ω be a domain in \mathbb{R}^n . Then the following Hardy-type inequality holds for all $u \in \dot{W}^{1,p}(\Omega)$, $p > 1$:*

$$\begin{aligned}
\int_{\Omega} |\nabla u|^p dx &\geq \frac{c_p \sqrt{\pi} \cdot \Gamma\left(\frac{n+p}{2}\right)}{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{n}{2}\right)} \left(\int_{\Omega} \int_{S^{n-1}} \frac{1}{\rho_\nu(x)^p} d\omega(\nu) |u(x)|^p dx \right. \\
&\quad \left. + (p-1) \left(\frac{|\mathbb{S}^{n-1}|}{n} \right)^{\frac{p}{n}} \int_{\Omega} \frac{|u(x)|^p}{|\Omega_x|^{\frac{p}{n}}} dx \right) \quad (2.2)
\end{aligned}$$

Proof. Clearly, we can assume $u \in C_0^\infty(\Omega)$. At first, we also assume that u is real valued. E.B.Davies arguments (see [6]) together with the one-dimensional inequality (Theorem 1.1) gives

$$\int_{\Omega} |\partial_\nu u|^p dx \geq c_p \int_{\Omega} \frac{|u(x)|^p}{\rho_\nu(x)^p} dx + c_p(p-1) \int_{\Omega} \left(\frac{2}{D_\nu(x)} \right)^p |u(x)|^p dx.$$

By definition, we have

$$|\partial_\nu u| = |\nu \cdot \nabla u| = |\nabla u| |\cos(\nu, \nabla u)|,$$

where $\cos(v, w)$ denotes the angle between $v, w \in \mathbb{R}^n$.

By inserting this into the above inequality and integrating both sides with respect to the normalized surface measure on \mathbb{S}^{n-1} , we get

$$\int_{\Omega} \int_{S^{n-1}} |\cos(\nu, \nabla u)|^p d\omega(\nu) |\nabla u(x)|^p dx \geq \quad (2.3)$$

$$c_p \left(\int_{\Omega} \int_{S^{n-1}} \left(\frac{1}{\rho_\nu(x)^p} + (p-1) \left(\frac{2}{D_\nu(x)} \right)^p \right) d\omega(\nu) |u(x)|^p dx \right). \quad (2.4)$$

Now note that

$$\int_{S^{n-1}} |\cos(\nu, \nabla u)|^p d\omega(\nu) = \int_{S^{n-1}} |\cos(e, \nu)|^p d\omega(\nu)$$

for any fixed unit vector $e \in \mathbb{R}^n$. Elementary calculations shows that

$$\int_{S^{n-1}} |\cos(e, \nu)|^p d\omega(\nu) = \frac{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{n}{2}\right)}{\sqrt{\pi} \cdot \Gamma\left(\frac{n+p}{2}\right)}.$$

By dividing both sides in (2.3), (2.4) with the latter quantity and using the above lemma, we get

$$\begin{aligned} \int_{\Omega} |\nabla u|^p dx &\geq \frac{c_p \sqrt{\pi} \cdot \Gamma\left(\frac{n+p}{2}\right)}{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{n}{2}\right)} \left(\int_{\Omega} \int_{S^{n-1}} \frac{1}{\rho_\nu(x)^p} d\omega(\nu) |u(x)|^p dx \right. \\ &\quad \left. + (p-1) \left(\frac{|\mathbb{S}^{n-1}|}{n} \right)^{\frac{p}{n}} \int_{\Omega} \frac{|u(x)|^p}{|\Omega_x|^{\frac{p}{n}}} dx \right) \end{aligned}$$

as desired. By standard density arguments, we get the same inequality for all real-valued $u \in \mathring{W}^{1,p}(\Omega)$.

Now take an arbitrary $v(x) \in C_0^\infty(\Omega)$ (not necessarily real-valued).

Then we have $|v| \in \mathring{W}^{1,p}(\Omega)$. Hence, we get the inequality (2.2) for the function $u(x) = |v(x)|$.

Since $|\nabla|v(x)|| \leq |\nabla v(x)|$ a.e (see for example E.H.Lieb and M.Loss [10], p.144), we get inequality (2.2) for all $u \in C_0^\infty(\Omega)$ and thus for all $u \in \mathring{W}^{1,p}(\Omega)$. This concludes the proof of the theorem. \square

For convex domains an easy geometric argument shows that

$$\int_{S^{n-1}} \frac{1}{\rho_\nu(x)^p} d\omega(\nu) \geq \int_{S^{n-1}} |\cos(e, \nu)|^p d\omega(\nu) \cdot \frac{1}{\delta(x)^p} = \frac{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{n}{2}\right)}{\sqrt{\pi} \cdot \Gamma\left(\frac{n+p}{2}\right) \delta(x)^p}.$$

For such domains we also know that $\Omega = \Omega_x$ for every $x \in \Omega$. Using the above theorem, we get

Theorem 2.2. *For any convex domain $\Omega \subset \mathbb{R}^n$ and $u \in \mathring{W}^{1,p}(\Omega)$ we have*

$$\int_{\Omega} |\nabla u|^p dx \geq c_p \int_{\Omega} \frac{|u(x)|^p}{\delta^p(x)} dx + \frac{a(p,n)}{|\Omega|^{\frac{p}{n}}} \int_{\Omega} |u(x)|^p dx,$$

where

$$a(p,n) = \frac{(p-1)^{p+1}}{p^p} \cdot \left(\frac{|\mathbb{S}^{n-1}|}{n} \right)^{\frac{p}{n}} \cdot \frac{\sqrt{\pi} \cdot \Gamma\left(\frac{n+p}{2}\right)}{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{n}{2}\right)}.$$

If Ω is not convex, we have the following counterpart of Corollary 3.1 in [9].

Corollary 2.1. *Suppose there exist a constant κ such that for each $y \in \partial\Omega$ and each $a > 0$ there exists a ball B with centre z disjoint from Ω and radius $\beta \geq \kappa a$, where $|z - y| = a$. Then there exists a constant $\theta \leq c_p$ such that*

$$\frac{c_p \cdot \sqrt{\pi} \cdot \Gamma\left(\frac{n+p}{2}\right)}{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{n}{2}\right)} \int_{S^{n-1}} \frac{1}{\rho_\nu(x)^p} d\omega(\nu) \geq \theta \frac{1}{\delta^p(x)}$$

and hence

$$\begin{aligned} \int_{\Omega} |\nabla u|^p dx &\geq \theta \int_{\Omega} \int_{S^{n-1}} \frac{1}{\rho_v(x)^p} d\omega(\nu) |u(x)|^p dx \\ &+ (p-1) \frac{c_p \sqrt{\pi} \cdot \Gamma\left(\frac{n+p}{2}\right)}{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{n}{2}\right)} \left(\frac{|\mathbb{S}^{n-1}|}{n}\right)^{\frac{p}{n}} \int_{\Omega} \frac{|u(x)|^p}{|\Omega_x|^{\frac{p}{n}}} dx. \end{aligned}$$

References

- [1] G.Barbatis, S.Filippas and A.Tertikas, ‘A unified approach to improved L^p Hardy inequalities with best constants’. Preprint, December 2000.
- [2] H.Brezis and M.Marcus, ‘Hardy’s inequalities revisited’, Dedicated to Ennio De Giorgi. *Ann.Scuola Norm. Sup.Pisa Cl.Sci* (4) vol. 25 (1997) no. 1–2 (1998), pp. 217–237.
- [3] E.B.Davies, ‘A review of Hardy inequalities’, *Oper. Theory Adv. Appl.*, vol. 110, (1998) pp. 55–67.
- [4] E.B.Davies, ‘Heat kernels and spectral theory’, [Cambridge, Cambridge University press, 1989].
- [5] E.B.Davies, ‘The Hardy constant’, *Quart.J.math. Oxford* (2), vol. 46, (1995) pp. 417–431.
- [6] E.B.Davies, ‘Spectral theory and differential operators’, *Cambridge studies in advanced mathematics* vol. 42, [Cambridge, Cambridge University Press, 1995].
- [7] G.Hardy, J.E.Littlewood and G.Pólya, ‘Inequalities, second edition’, *Cambridge Mathematical Library*, [Cambridge, Cambridge University Press, 1952].
- [8] G.Hardy, ‘Note on a theorem of Hilbert’, *Math. Zeitschr.*(6),(1920) pp. 314–317.
- [9] M.Hoffmann–Ostenhof, T.Hoffmann–Ostenhof and A.Laptev, ‘A geometrical version of Hardy’s inequality’. Accepted by *J.Func.Anal.*
- [10] E.H.Lieb and M.Loss, ‘Analysis’, *Graduate Studies in Mathematics* vol. 14, [USA, American Mathematical Society 1997].
- [11] T.Matskewich and P.E.Sobolevskii, ‘The best possible constant in a generalized Hardy’s inequality for convex domains in \mathbb{R}^n ’, *Nonlinear Anal.* vol. 28 (1997) pp. 1601–1610.
- [12] V.G.Maz’ja, ‘Sobolev Spaces’, [Berlin, Springer–Verlag 1985].
- [13] B.Opic and A.Kufner, ‘Hardy–type inequalities’, *Pitman Research notes in mathematics series* vol. 219, [London, Longman Group UK Limited, 1990].

Jesper Tidblom, Ph.D. student. Department of Mathematics, University of Stockholm, 106 91 Stockholm, Sweden.

E–mail : jespert@matematik.su.se