ISSN: 1401-5617

Koszul homology and Lie algebras with application to generic forms and points

Ralf Fröberg
Clas Löfwall

Research Reports in Mathematics
Number 8, 2000

Department of Mathematics
Stockholm University

Electronic versions of this document are available at
http://www.matematik.su.se/reports/2000/8

Date of publication: June 2, 2000
1991 Mathematics Subject Classification: Primary 13D07 Secondary 14M99, 16E45.

Postal address:
Department of Mathematics
Stockholm University
S-106 91 Stockholm
Sweden

Electronic addresses:
http://www.matematik.su.se
info@matematik.su.se

Koszul homology and Lie algebras with application to generic forms and points

R. Fröberg* C. Löfwall†

27 maj 2000

1 Introduction

Consider s generic points in projective space P_k^n. The Hilbert series of the corresponding coordinate ring $A = S/I = k[x_0, \dots, x_n]/I$ is known to be $\sum_{i\geq 0} \min(\binom{n+i}{i}, s)z^i$ [Ge-Or 81]. The Koszul homology of A, $H = \mathrm{Tor}^S(A,k)$, is known to satisfy $H_{i,j} = 0$ if $j \neq d+i-1, d+i$ for a certain d (depending on n and s), but there are a lot of unanswered questions. The Minimal Resolution Conjecture (MRC) states that for all i and all j, $H_{i,j} = 0$ or $H_{i+1,j} = 0$, and the special case that this is true for $i = 1$ is called the Ideal Generating Conjecture (IGC). The MRC has recently been disproved [Ei-Po 99], while IGC is still open. In this paper we will introduce a new approach to study these problems, namely through the "homotopy Lie algebra", $\mathfrak{g}_A$, of A.

The MRC may be reformulated in terms of $\mathfrak{g}_A$. In fact, there are certain numbers m and t (depending on n and s) such that if $d > 2$ then MRC holds if and only if $\dim_k \mathfrak{g}_A^{m+1,m+d-1} = t$ and $\mathfrak{g}_A^{m+2,m+d} = 0$. When $d = 2$ the situation is a bit more complicated. Let η_A be the "diagonal" Lie algebra of $\mathfrak{g}_A$; i.e., $\eta_A^i = \mathfrak{g}_A^{i,i}$ and let $L = \oplus_{i\geq 2}\eta_A^i$ and finally $Q^i = (L/[L,L])^i$. Then if $d = 2$, MRC holds if and only if $\dim_k Q^{m+1} = t$ and $Q^{m+2} = 0$.

Even if the case $d > 2$ is easier, we do not know any application of it. However in the case when $d = 2$, the Lie algebra η_A is explicitly given as the free Lie algebra on $n+1$ odd variables modulo the ideal generated by the squares of s generic linear forms. We have used a Mathematica program called "liedim" and its C-version "cbas" to compute the space Q above for some values of n and s. In this way we have been able to prove MRC and IGC in some new cases.

We have also studied ideals generated by generic quadratic forms in polynomial rings and exterior algebras and ideals generated by generic quadratic Lie elements in free Lie algebras.

The paper is organized as follows. In Section 2 the Koszul dual is defined for general superalgebras, in particular for graded commutative algebras. This will be applied to the Koszul homology algebra, which has generators of both even and odd degrees. In Section 3 the homotopy Lie algebra is introduced for graded commutative algebras through minimal models and in Section 4 a theorem is

* *email* ralff@matematik.su.se
† *email* clas@matematik.su.se

proved which relates parts of the Koszul homology algebra and the homotopy Lie algebra by means of Koszul duality. In fact, since we are dealing with superalgebras, our theorem may also be applied to quotients of exterior algebras (or to quotients of tensor products of exterior algebras and polynomial algebras). However, in the exterior algebra case, $\mathrm{Tor}^S(A,k)$ is in general infinite-dimensional and the corresponding Lie algebra is an "ordinary" Lie algebra (non-super).

In Section 5 we introduce the MRC and IGC conjectures and apply the theorem in Section 4 in order to reformulate the conjectures in terms of the Lie algebra L associated to A. We also prove a "monotonicity" property, which implies that if counterexamples to MRC exists for a given n, then there has to be counterexamples for special values of s. For IGC this holds for at most two values of s for each n. In Section 6 we present some theorems about MRC and IGC obtained by computations and by the theory from the previous section. The computations are presented in Section 11.

In Section 7 we study rings of the form $k[x_1,\dots,x_n]/(f_1,\dots,f_r)$, where $f_1,\dots,f_r$ are generic quadratic forms. We prove that the ring is Koszul if and only if $r \le n$ or $r \ge \binom{n+1}{2} - n^2/4$. The corresponding result for Lie algebras is also obtained. We study the series of Lie algebras obtained from generic quadratic algebras or defined by generic quadratic Lie relations. We make some conjectures and prove them in some special cases. In Section 8 we state a conjecture about the Lie algebra for the coordinate ring of a set of generic points, and we prove it in some cases. In Section 9 we study the Poincaré series for these coordinate rings. In Section 10 finally, we study the quotient of exterior algebras with generic quadratic forms and their corresponding (ordinary) Lie algebras. In particular we study the exterior algebra in 5 variables modulo 2 or 3 generic quadratic forms. The Hilbert series turned out to be not the ones which were expected.

2 The Koszul dual

We will consider $\mathbb{Z}_2\times\mathbb{N}$-graded (or $\mathbb{Z}_2\times\mathbb{N}\times\mathbb{N}$-graded) associative algebras or Lie algebras over a field k of characteristic different from two. The $\mathbb{Z}_2$-grading (also called the "parity") of an element x is denoted $|x|$ and we ususally talk about "even" and "odd" elements corresponding to whether $|x|=0$ or $|x|=1$. This grading makes our algebras "superalgebras", and it is this grading that determines the sign when two elements in a formula are interchanged; e.g., the graded commutator is defined as $[a,b]=ab-(-1)^{|a||b|}ba$ and the graded Jacobi identity is $[[a,b],c]=[a,[b,c]]+(-1)^{|b||c|}[[a,c],b]$. The additional $\mathbb{N}$-gradings are called "weights" and they have no effect on signs in formulas.

The Koszul dual $A^!$ of a $\mathbb{Z}_2\times\mathbb{N}$-graded connected k-algebra $A=k\oplus A_1\oplus A_2\oplus\cdots$ is again a $\mathbb{Z}_2\times\mathbb{N}$-graded connected k-algebra and may be defined as the subalgebra of $\mathrm{Ext}_A(k,k)$ generated by the elements of homological degree and weight equal to one. The algebra A is called a Koszul algebra if $A^!=\mathrm{Ext}_A(k,k)$. The Koszul dual may be computed using the cobar construction $(\mathrm{T}((sA_+)^*),d)$, cf [Ad 60]. Here A_+ is the set of elements of positive weight, which is considered to be concentrated in homological degree zero, T stands for "tensor algebra" and s is the suspension functor, which changes parity and raises the homological degree by one, while the weight is left unchanged by s. The differential d is the derivation which extends the map

$$(sA_+)^* \overset{s}{\to} A_+^* \overset{m^*}{\to} (A_+ \otimes A_+)^* \overset{(\phi_{A_+})^{-1}}{\to} (sA_+)^* \otimes (sA_+)^* ,$$

where $m\colon A_+ \otimes A_+ \to A_+$ is multiplication and ϕ_{A_+} is a natural isomorphism ϕ_V applied to $V = A_+$. It is defined as follows. Let V be a $\mathbb{Z}_2 \times \mathbb{N}$-graded vector space, then $\phi_V\colon (sV)^* \otimes (sV)^* \to (V \otimes V)^*$ is defined by $\phi(f \otimes g) = (f \otimes g) \circ (s \otimes s)$ for $f, g \in (sV)^*$. Following the strict sign rule, this means that the following holds for $a, b \in V$.

$$(\phi(f \otimes g))(a \otimes b) = (-1)^{|a|+|g||sa|} f(sa)g(sb) = (-1)^{|a|+|f||g|} f(sa)g(sb) = -(-1)^{|sa||b|} f(sa)g(sb) .$$

It is easy to prove (cf [Pr 70], [Lö 86]) that

$$A^! = \mathrm{T}((sA_1)^*)/\langle d((sA_2)^*)\rangle .$$

Here $A^!$ only depends on the "1-2-algebra of A",

$$\mathrm{T}(A_1)/\langle \ker(A_1 \otimes A_1 \to A_2)\rangle ,$$

which in fact is equal to $(A^!)^!$.

This gives the following description of the Koszul dual. Let V be a $\mathbb{Z}_2$-graded finite-dimensional (or locally finite-dimensional $\mathbb{Z}_2 \times \mathbb{N}$-graded) vector space and R a homogeneous subspace of $V \otimes V$. Then

$$(\mathrm{T}(V)/\langle R\rangle)^! = \mathrm{T}((sV)^*)/\langle \phi_V^{-1} R^\perp\rangle ,$$

where s is the parity switcher, $R^\perp = \{f \in (V \otimes V)^*;\ f(R) = 0\}$ and $\phi_V\colon\ (sV)^* \otimes (sV)^* \to (V \otimes V)^*$ is the isomorphism defined above. In coordinates this means the following (we have changed the formulas by a global -1). If $\{x, y, \dots\}$ is a basis for V and $\{X, Y, \dots\}$ is a basis for $(sV)^*$ dual to $\{sx, sy, \dots\}$, then an element $X \otimes Y$ is considered to operate on a basis element $a \otimes b$ by the rule $(X \otimes Y)(a \otimes b) = 0$ if $a \neq x$ or $b \neq y$, $(X \otimes X)(x \otimes x) = 1$ and

$$(X \otimes Y)(x \otimes y) = \begin{cases} -1 & \text{if } x \text{ is even and } y \text{ is odd} \\ 1 & \text{otherwise.} \end{cases}$$

Observe that this somewhat strange rule follows from the fact that the suspension s has to pass x before Y passes sx.

Example 2.1 Let $|x| = 0$, $|y| = 1$. Then

$$(k\langle x, y\rangle/\langle xy - yx, y^2\rangle)^! = k\langle X, Y\rangle/\langle XY - YX, X^2\rangle ,$$

where $|X| = 1$, $|Y| = 0$.

This example is in accordance with the known fact that the Ext-algebra of a free graded commutative algebra is free graded commutative with a parity switch (cf. [Ma 88] 3.10) . See Proposition 2.2 below.

For a $\mathbb{Z}_2$-graded vector space V, let $[V,V]_{\mathcal{F}}$ denote the subspace of $V\otimes V$ generated by the graded commutators. The subscript $\mathcal{F}$ refers to the fact that $[V,V]_{\mathcal{F}}$ is contained in $\mathcal{F}(V)$, the free Lie algebra on V, considered as the Lie subalgebra generated by V in the free associative algebra $\mathrm{T}(V)$. Then $\mathrm{U}(\mathcal{F}(V)) = \mathrm{T}(V)$, where U stands for "universal enveloping algebra". The free graded commutative algebra on V, denoted $\bigwedge(V)$, is defined as $\mathrm{T}(V)/\langle [V,V]_{\mathcal{F}}\rangle$. Now $V\wedge V = (V\otimes V)/[V,V]_{\mathcal{F}}$ and hence $(V\wedge V)^*$ may (and will) be identified with $([V,V]_{\mathcal{F}})^{\perp}\subset (V\otimes V)^*$.

Proposition 2.2 *Let V be a $\mathbb{Z}_2\times\mathbb{N}$-graded locally finite-dimensional vector space and let s be the parity switcher. Let R be a subspace of $V\otimes V$ which contains $[V,V]_{\mathcal{F}}$ and let $\overline{R} = R/[V,V]_{\mathcal{F}}$. Consider the map $\phi_V = \phi\colon (sV)^*\otimes (sV)^* \to (V\otimes V)^*$ defined (as above) by $\phi(f\otimes g)(a\otimes b) = (-1)^{|a|+|g||sa|}f(sa)g(sb)$. Then*

$$
\begin{aligned}
&(i) && \phi[(sV)^*,(sV)^*]_{\mathcal{F}} = ([V,V]_{\mathcal{F}})^{\perp}\\
&(ii) && (\textstyle\bigwedge(V)/\langle\overline{R}\rangle)^! \cong \mathrm{U}(\mathcal{F}((sV)^*)/\langle\phi^{-1}R^{\perp}\rangle)\\
&(iii) && (\textstyle\bigwedge(V))^! \cong \bigwedge((sV)^*)\\
&(iv) && \overline{R}^* \cong [(sV)^*,(sV)^*]_{\mathcal{F}}/\phi^{-1}R^{\perp}.
\end{aligned}
$$

Proof. (i). Let $f,g\in(sV)^*$ and $a,b\in V$. Then

$$
\begin{aligned}
&\phi(f\otimes g)(a\otimes b - (-1)^{|a||b|}b\otimes a) =\\
&\qquad (-1)^{|a||b|+|b|+1}f(sa)g(sb) - (-1)^{|a|+1}f(sb)g(sa) \quad \text{and}\\
&\phi(g\otimes f)(a\otimes b - (-1)^{|a||b|}b\otimes a) =\\
&\qquad (-1)^{|f||g|+|a|}g(sa)f(sb) - (-1)^{|f||g|+|a||b|+|b|}g(sb)f(sa).
\end{aligned}
$$

Hence $\phi(f\otimes g) - (-1)^{|f||g|}\phi(g\otimes f)$ is zero on $a\otimes b - (-1)^{|a||b|}b\otimes a$ and hence $\phi[(sV)^*,(sV)^*]_{\mathcal{F}}\subset([V,V]_{\mathcal{F}})^{\perp}$. The other inclusion follows by dimension reasons.

(ii). This follows directly from (i) and the description of the Koszul dual above, since $\bigwedge(V)/\langle\overline{R}\rangle \cong \mathrm{T}(V)/\langle R\rangle$ and

$$
\mathrm{T}((sV)^*)/\langle\phi^{-1}R^{\perp}\rangle \cong \mathrm{U}\left(\mathcal{F}((sV)^*)/\langle\phi^{-1}R^{\perp}\rangle\right),
$$

which is true since, $\phi^{-1}R^{\perp}\subset\phi^{-1}([V,V]_{\mathcal{F}})^{\perp} = [(sV)^*,(sV)^*]_{\mathcal{F}}$.

(iii). This follows from (i) and (ii) with $R = [V,V]_{\mathcal{F}}$.

(iv). Using (i) we get

$$
\begin{aligned}
\overline{R}^* \;=\;& (R/[V,V]_{\mathcal{F}})^* \cong ([V,V]_{\mathcal{F}})^{\perp}/R^{\perp} \cong\\
& \phi^{-1}([V,V]_{\mathcal{F}})^{\perp}/\phi^{-1}R^{\perp} = [(sV)^*,(sV)^*]_{\mathcal{F}}/\phi^{-1}R^{\perp}.
\end{aligned}
$$

□

In general, for an augmented k-algebra A, the subalgebra of $\mathrm{Ext}_A(k,k)$ generated by $\mathrm{Ext}^1_A(k,k)$ has relations of degree greater than two. However the relations of degree two may be described explicitly.

Proposition 2.3 *Let A be an augmented k-algebra with augmentation ideal I and let $\mathrm{gr}(A) = k \oplus I/I^2 \oplus I^2/I^3 \oplus \cdots$ be the graded associated k-algebra. Then the 1-2-algebra of $\mathrm{Ext}_A(k,k)$ is isomorphic to the Koszul dual of $\mathrm{gr}(A)$; i.e.,*

$$(\mathrm{Ext}_A(k,k)^!)^! \cong (\mathrm{gr}(A))^! .$$

Proof. All we need is to consider the beginning of the cobar construction, $(sI)^* \xrightarrow{d} (sI)^* \otimes (sI)^*$, where d is defined as above. We have $\mathrm{Ext}^1_A(k,k) = (sI^2)^\perp$. Hence

$$\ker\left(\mathrm{Ext}^1_A(k,k) \otimes \mathrm{Ext}^1_A(k,k) \to \mathrm{Ext}^2_A(k,k)\right) = \mathrm{im}(d) \cap \left((sI^2)^\perp \otimes (sI^2)^\perp\right) .$$

With ϕ and m as above we have $\phi(\mathrm{im}(d)) = (\ker(m))^\perp$ and

$$\phi\left((sI^2)^\perp \otimes (sI^2)^\perp\right) = (I^2 \otimes I + I \otimes I^2)^\perp .$$

Hence

$$\phi \ker\left(\mathrm{Ext}^1_A(k,k) \otimes \mathrm{Ext}^1_A(k,k) \to \mathrm{Ext}^2_A(k,k)\right) = \left(I^2 \otimes I + I \otimes I^2 + \ker(m)\right)^\perp$$

which equals $(m^{-1}(I^3))^\perp$, since $m(I^2 \otimes I) = I^3$. Thus the 1-2-algebra of $\mathrm{Ext}_A(k,k)$ is isomorphic to $\mathrm{T}((sI^2)^\perp)/\langle \phi^{-1}(m^{-1}(I^3))^\perp \rangle$.

On the other hand

$$(\mathrm{gr}(A))^! = \mathrm{T}((sI/I^2)^*)/\langle \phi^{-1}_{I/I^2}(\ker(I/I^2 \otimes I/I^2 \to I^2/I^3))^\perp \rangle$$

Looking at $(I/I^2 \otimes I/I^2)^*$ as a subspace of $(I \otimes I)^*$, the space $\left(\ker(I/I^2 \otimes I/I^2 \to I^2/I^3)\right)^\perp$ will be identified with $(m^{-1}(I^3))^\perp$. Hence

$$(\mathrm{gr}(A))^! \cong \mathrm{T}((sI^2)^\perp)/\langle \phi_I^{-1}(m^{-1}(I^3))^\perp \rangle .$$

□

We end this section with a proposition which, combined with Proposition 2.3, will be used in Section 4.

Proposition 2.4 *Let L be a $\mathbb{Z}_2 \times \mathbb{N}^+$-graded Lie algebra, locally finite-dimensional over k. Let $\mathrm{gr}(L)$ denote the graded associated Lie algebra with respect to the filtration $L \supset [L,L] \supset [L,[L,L] \supset \cdots$ and let $\mathrm{gr}(\mathrm{U}(L))$ denote the graded associated algebra with respect to the filtration obtained from powers of the augmentation ideal. Then $\mathrm{gr}(\mathrm{U}(L))$ and $\mathrm{U}(\mathrm{gr}(L))$ are naturally isomorphic as graded algebras.*

Proof. There is a natural map of Lie algebras from $\mathrm{gr}(L)$ to the Lie algebra associated to the algebra $\mathrm{gr}(\mathrm{U}(L))$ and hence there is induced a natural map of graded algebras $\mathrm{U}(\mathrm{gr}(L)) \to \mathrm{gr}(\mathrm{U}(L))$. This map is surjective, since both algebras are generated by $L/[L,L]$ (depending on the assumption that L has a positive grading). Since, by assumption, both algebras are locally finite-dimensional, it follows that the map is an isomorphism, since the algebras have the same Hilbert series.

□

3 The homotopy Lie algebra

Let V be a $\mathbb{Z}_2 \times \mathbb{N}^+$-graded vector space over k with basis $\{x_1, \dots, x_n\}$. We will also denote the free graded commutative algebra on V, $\bigwedge(V)$, by $k[x_1, \dots, x_n]$. It is the free graded associative algebra $k\langle x_1, \dots, x_n\rangle$ modulo the graded commutators $x_i x_j - (-1)^{|x_i||x_j|} x_j x_i$. Sometimes V will be infinite-dimensional, but always finite-dimensional in each weight.

We will consider k-algebras $A = S/I$, where $S = k[x_1, \dots, x_n]$ and I is a homogeneous ideal with respect to both parity and weight. To such a k-algebra A, we will define a $\mathbb{Z}_2 \times \mathbb{N}^+ \times \mathbb{N}^+$-graded Lie algebra $\mathfrak{g}_A$ – the homotopy Lie algebra of A. It may be defined by the property $U(\mathfrak{g}_A) = \mathrm{Ext}_A(k,k)$, but we will give a more explicit definition (and more suitable for our purposes) using the minimal model of A (cf [Lö 94]).

The minimal model of A is a differential algebra $(\bigwedge V, d)$, where V is a $\mathbb{Z}_2 \times \mathbb{N} \times \mathbb{N}^+$-graded vector space and where the first degree is the parity, the second degree is the homological degree and the third degree is the weight. The differential d is a derivation which changes parity, lowers the homological degree by one and preserves weight. Moreover d maps V to $\bigwedge^{\geq 2} V = V \wedge V + V \wedge V \wedge V + \cdots$. The component of d mapping V to $V \wedge V$ is called d_2 (its extension as a derivation also satisfies $d_2^2 = 0$).

The ring A is considered as a differential algebra with zero differential and concentrated in homological degree zero. There is a surjective map $\epsilon\colon (\bigwedge V, d) \to A$ preserving all degrees and inducing an isomorphism in homology. This means that $H_i(\bigwedge V, d) = 0$ for homological degrees $i > 0$ and if $A = k[x_1, \dots, x_n]/I$ then V in homological degree zero has a basis $X_1, \dots, X_n$, $\epsilon(X_i) = x_i$ and $\epsilon(\mathrm{im}(d)) = I$. The construction of $(\bigwedge V, d)$ is a straightforward procedure of killing cycles in a minimal way (see [Lö 94] for more details).

The Lie algebra $\mathfrak{g}_A$ is defined by $(sV)^*$ as a vector space, where s changes parity, raises homological degree by one and preserves weight. The Lie product on $(sV)^*$ is in principle defined as the dual of d_2. More precisely, it is defined as a map $[(sV)^*, (sV)^*]_{\mathcal{F}} \to (sV)^*$ in the following way (cf. Proposition 2.2),

$$[(sV)^*, (sV)^*]_{\mathcal{F}} \xrightarrow{\phi} ([V,V]_{\mathcal{F}})^{\perp} = (V \wedge V)^* \xrightarrow{d_2^*} V^* \xrightarrow{s^{-1}} (sV)^* .$$

This map preserves all degrees and it may be checked that it satisfies the graded Jacobi identity. In coordinates the Lie product may be given as follows (cf [Av 84], [Ha 92]).

Suppose $\{x, y, z, \dots\}$ is an ordered basis for V and let $\{X, Y, Z, \dots\}$ denote the basis for $(sV)^*$ dual to $\{sx, sy, sz, \dots\}$. If $d_2 x = \sum_{y \leq z} c_{xyz} yz$ then,

$$\begin{aligned} [Z,Y] &= -\textstyle\sum_x (-1)^{|y|} c_{xyz} X \qquad \text{if } y < z \\ (Y)^2 &= -\textstyle\sum_x c_{xyy} X \qquad \text{if } Y \text{ is odd.} \end{aligned}$$

Inside $\mathfrak{g}_A$ there is the Lie subalgebra η_A generated by $\mathfrak{g}_A^{1,1}$ (homological degree = weight = 1). It is proved in [Lö 94] that η_A coincides with the diagonal subalgebra $\oplus_{p \geq 1}^{\infty} \mathfrak{g}_A^{p,p}$ and it is proved in [Pr 70], [Lö 86] that $\mathrm{U}(\eta_A) = A^!$.

Definition 3.1 *Let $\mathfrak{g}_A$ be the homotopy Lie algebra of A and η_A the Lie subalgebra of $\mathfrak{g}_A$ generated by $\mathfrak{g}_A^{1,1}$. The Lie algebra $L = (\eta_A)^{\geq 2} = [\eta_A, \eta_A]$ is called the "Lie algebra associated to A". It has two degrees, a parity and a weight (≥ 2).*

Let L be the Lie algebra associated to A and let $\mathrm{gr}(L)$ denote the graded associated Lie algebra obtained from the lower central series $L \supset [L,L] \supset [L,[L,L]] \supset \cdots$; i.e.,

$$\mathrm{gr}(L) = (\mathrm{gr}(L))_1 \oplus (\mathrm{gr}(L))_2 \oplus \cdots = L/[L,L] \oplus [L,L]/[L,[L,L]] \oplus \cdots .$$

Each $(\mathrm{gr}(L))_i$ has an induced grading from L, $(\mathrm{gr}(L))_i = \oplus_j (\mathrm{gr}(L))_i^j$. The following theorem relates $\mathrm{gr}(L)$ to $\mathrm{Tor}^S(A,k)$, which is the homology of the Koszul complex of A if x_i is even for all i. The theorem may be seen as a study of an edge in Avramov's spectral sequence ([Av 74]), which has as E^2-term $\mathrm{Tor}^{\mathrm{Tor}^S(A,k)}(k,k)$ and which converges to $\mathrm{U}(\mathfrak{g}_A^{\geq 2})^*$. At this point we want to thank J.-E. Roos, who conjectured part one of the theorem below and encouraged us to supply a proof.

4 The Koszul homology in terms of the homotopy Lie algebra

Theorem 4.1 *Let $A = S/I = k[x_1, \ldots, x_n]/I$ be a $\mathbb{Z}_2 \times \mathbb{N}$-graded connected k-algebra, commutative in the graded sense by the first degree, which is called the "parity"; the second degree is called the "weight". Let $H = \oplus H_{i,j} = \oplus \mathrm{Tor}_{i,j}^S(A,k)$ (here i is the homological degree and j is the weight), which is considered as a graded algebra in the following way; $H = k \oplus H_{(1)} \oplus H_{(2)} \oplus \cdots$, where $H_{(i)} = \oplus_j H_{j-i,j}$. Furthermore parity and weight in A defines a bigrading on each $H_{(i)}$ (which is compatible with the multiplication in H). Let L be the Lie algebra associated to A and $\mathrm{gr}(L)$ the graded associated Lie algebra with respect to the lower central series. Then $\mathrm{gr}(L)$ has, besides a parity, two gradings, $\mathrm{gr}(L) = \oplus_{i \geq 1, j \geq 2} (\mathrm{gr}(L))_i^j$ (see above). The following natural isomorphisms hold, where in (ii) the operator s just changes the parity.*

- (*i*) *The 1-2-algebra of H is isomorphic to the Koszul dual of* $\mathrm{U}(\mathrm{gr}(L))$*; i.e.,* $(H^!)^! \cong (\mathrm{U}(\mathrm{gr}(L)))^!$
- (*ii*) $H_{(1)} \cong s(L/[L,L])^*$
- (*iii*) $\ker(H_{(1)} \wedge H_{(1)} \to H_{(2)}) \cong ([L,L]/[L,[L,L]])^*$
- (*iv*) *If A is Koszul, then $H \cong \mathrm{Ext}_{\mathrm{U}(L)}(k,k)$ as algebras, in particular* $H_{j-i,j} = H_{(i),j} \cong (\mathrm{Ext}^i_{\mathrm{U}(L)}(k,k))_j$.

Corollary 4.2 *With notation as in the theorem, suppose x_i is even for all i. Then*

- (*i*) $L^j = [L,L]^j$ *for* $j \geq n+2$
- (*ii*) *If A is Koszul, then* $\mathrm{gldim}(\mathrm{U}(L)) < \infty$.

Corollary 4.3 *With notation as in the theorem, we have*

$$
\begin{aligned}
&(i) && \dim_k(H_{1,2})_{even} = \dim_k(L^2)_{odd} \\
&(ii) && \dim_k(H_{1,2})_{odd} = \dim_k(L^2)_{even} \\
&(iii) && \dim_k H_{2,3} = \dim_k L^3 \\
&(iv) && \dim_k H_{3,4} = \dim_k L^4/[L^2, L^2] \\
&(v) && \dim_k(\ker(H_{1,2} \wedge H_{1,2} \to H_{2,4})) = \dim_k[L^2, L^2] \\
&(vi) && \dim_k(\ker(H_{1,2} \otimes H_{2,3} \to H_{3,5}))_{even} = \dim_k([L^2, L^3])_{even} \\
&(vii) && \dim_k(\ker(H_{1,2} \otimes H_{2,3} \to H_{3,5}))_{odd} = \dim_k([L^2, L^3])_{odd} \\
&(viii) && \dim_k \ker\left(H_{2,3} \wedge H_{2,3} \oplus H_{1,2} \otimes H_{3,4} \to H_{4,6}\right) = \\
& && \qquad \dim_k \left([L^3, L^3] + [L^2, L^4]\right) / [L^2, [L^2, L^2]] \,.
\end{aligned}
$$

Remark 4.4 It has been known since long that $\dim_k H_{2,3} = \dim_k L^3$ and this was the starting point for our study of generic points and it is mainly this fact we will use in the applications of the theorem.

Proof of Theorem 4.1. We will give three different proofs. Even if the second one is short, we have included the other two, since we believe the methods used there are interesting in themselves. The first two use the minimal model of A. The third uses the cobar construction of (the dual of) the Koszul complex and the perturbation theory developed by Gugenheim, Stasheff, Lambe and others [Gu-La-St 91], [Hu-Ka 91], [Jo-La 00].

First proof. Let $(\bigwedge(V), d)$ be a minimal model of A as described above. In particular this is a free resolution of A as a module over $S \cong \bigwedge(V_0)$. Hence $\mathrm{Tor}^S(A, k)$ may be computed as the homology of $\bigwedge(V) \otimes_S k = \bigwedge(V_{\geq 1})$ with the induced differential $\overline{d}$. Due to the minimality of $(\bigwedge(V), d)$ and the fact that the generators in S have positive weight, we may conclude that the weight is always greater than the homological degree. Let W be the subspace of $V_{\geq 1}$ defined by elements of minimal possible weight; i.e., $W = \oplus_{i \geq 1} V_{i,i+1}$. Then $L = (sW)^*$. By degree reasons, the differential $\overline{d}$ maps W to $W \wedge W$ and we will call this map d_2. Hence $(\bigwedge(W), d_2)$ is a subcomplex of $(\bigwedge(V_{\geq 1}), \overline{d})$ and moreover $(\overline{d})^{-1}(W \wedge W) = W$. Put $W^{(1)} = \ker(d_2)$ and $W^{(2)} = d_2^{-1}(W^{(1)} \wedge W^{(1)})$. Then $W^{(1)} \subset W^{(2)} \subset W$. We have

$$
\begin{aligned}
H_{(1)} &= W^{(1)} \\
\ker(H_{(1)} \wedge H_{(1)} \to H_{(2)}) &= d_2(W^{(2)}) \,.
\end{aligned}
$$

Claim.

$$
\begin{aligned}
(sW^{(1)})^\perp &= [L, L] \\
(sW^{(2)})^\perp &= [L, [L, L]]
\end{aligned}
$$

Once this is proved, we get $(sH_{(1)})^* = (sW^{(1)})^* = (sW)^*/(sW^{(1)})^\perp = L/[L, L]$ which proves (ii). We also get $(sW^{(2)})^* = L/[L, [L, L]]$. Let d_{22} denote the restriction of d_2 to $W^{(2)}$, then $d_{22} \colon W^{(2)} \to W^{(1)} \wedge W^{(1)}$. By the above $\ker(H_{(1)} \wedge H_{(1)} \to H_{(2)}) = \mathrm{im}(d_{22})$. To prove (i) we have to prove that $\phi^{-1}(\mathrm{im}(d_{22}))^\perp$ is equal to the kernel of

$$
[L/[L, L], L/[L, L]]_{\mathcal{F}} \to [L, L]/[L, [L, L]] \,.
$$

But this is given as the kernel of the composition of the following maps.

$$[(sW^{(1)})^*, (sW^{(1)})^*]_{\mathcal{F}} \xrightarrow{\phi} (W^{(1)} \wedge W^{(1)})^* \xrightarrow{d_{22}^*} (W^{(2)})^* \to (sW^{(2)})^*$$

Since the first and the last map are isomorphisms, this kernel is $\phi^{-1}\ker(d_{22}^*) = \phi^{-1}(\mathrm{im}(d_{22}))^{\perp}$ which proves (i).

Now (iii) follows from Proposition 2.2 (iv).

If A is Koszul then $W = V_{\geq 1}$ and hence $(\bigwedge(V_{\geq 1}), \overline{d})$ is the same as $(\bigwedge(W), d_2)$, which is the (generalized) standard Chevalley-Eilenberg complex (cf [Ch-Ei 48]) to compute the cohomology of the Lie algebra $(sW)^*$ (here the "wedge-degree" is the cohomological degree and the second degree of W is the weight). Hence $H = \mathrm{H}((\bigwedge(W), d_2)) = \mathrm{Ext}_{\mathrm{U}(L)}(k,k)$ which proves (iv).

Proof of Claim.

We have that $[L,L]$ is the image of the map

$$[(sW)^*, (sW)^*]_{\mathcal{F}} \xrightarrow{\phi} (W \wedge W)^* \xrightarrow{d_2^*} W^* \to (sW)^*.$$

Hence $[L,L] = s^{-1}(\mathrm{im}(d_2^*)) = s^{-1}(\ker(d_2))^{\perp} = (sW^{(1)})^{\perp}$.

To prove the second claim, we have to prove that the map above maps the subspace $[(sW)^*, (sW^{(1)})^{\perp}]_{\mathcal{F}}$ onto $(sW^{(2)})^{\perp}$. We will do this in two steps

1. $\phi([(sW)^*, (sW^{(1)})^{\perp}]_{\mathcal{F}}) = (W^{(1)} \wedge W^{(1)})^{\perp}$
2. $d_2^*((W^{(1)} \wedge W^{(1)})^{\perp}) = (W^{(2)})^{\perp}$.

To prove 1., we first observe the following general fact. Suppose $B \to C$ is a surjective map with kernel D. Then $\mathrm{T}(B)/\langle D \rangle \cong \mathrm{T}(C)$ (use right exactness of the tensor product, diagram chasing and induction). Hence also $\mathcal{F}(B)/\langle D \rangle \cong \mathcal{F}(C)$. In particular $\ker([B,B]_{\mathcal{F}(B)} \to [C,C]_{\mathcal{F}(C)}) = [B,D]_{\mathcal{F}(B)}$. Now since ϕ is a natural isomorphism, ϕ induces an isomorphism

$$\ker\left\{[(sW)^*, (sW)^*]_{\mathcal{F}} \to [(sW^{(1)})^*, (sW^{(1)})^*]_{\mathcal{F}}\right\} \to$$
$$\ker\left\{(W \wedge W)^* \to (W^{(1)} \wedge W^{(1)})^*\right\}.$$

By the general fact above, the first kernel is $[(sW)^*, (sW^{(1)})^{\perp}]_{\mathcal{F}}$. Since the second kernel is $(W^{(1)} \wedge W^{(1)})^{\perp}$, the first statement is proved. The second statement is a direct consequence of the following general fact (which is easily proven). Suppose $f\colon B \to C$ is a map of vector spaces and let D be a subspace of C. Then $f^*(D^{\perp}) = (f^{-1}(D))^{\perp}$.

Second proof.

We will use the same constructions and notations as in the first proof. Even if A is not Koszul, we may argue in the same way as in the proof of (iv) above to conclude that $\mathrm{H}(\bigwedge(W), d_2) = \mathrm{Ext}_{\mathrm{U}(L)}(k,k)$. Since $(\bigwedge(W), d_2)$ is a subcomplex of $(\bigwedge(V_{\geq 1}), \overline{d})$ we get a map of algebras $\mathrm{Ext}_{\mathrm{U}(L)}(k,k) \to H$. We will study this map in low degrees and prove that the 1-2-algebra of $\mathrm{Ext}_{\mathrm{U}(L)}(k,k)$ is isomorphic to the 1-2-algebra of H. This, combined with Proposition 2.3 and Proposition 2.4, proves (i). Then (ii) follows from (i) and (iii) and (iv) are proved as before.

Obviously we have $\mathrm{Ext}^1_{\mathrm{U}(L)}(k,k) = W^{(1)} = H_{(1)}$.

Since $(\overline{d})^{-1}(W \wedge W) = W$, we have

$$\mathrm{im}(d_2) \cap (W^{(1)} \wedge W^{(1)}) = \mathrm{im}(\overline{d}) \cap (W^{(1)} \wedge W^{(1)}) .$$

Hence,

$$\begin{aligned} (\mathrm{Ext}^1_{\mathrm{U}(L)}(k,k))^2 &= W^{(1)} \wedge W^{(1)}/\mathrm{im}(d_2) \cap (W^{(1)} \wedge W^{(1)}) \\ &= W^{(1)} \wedge W^{(1)}/\mathrm{im}(\overline{d}) \cap (W^{(1)} \wedge W^{(1)}) = (H_{(1)})^2 . \end{aligned}$$

It follows that

$$\ker \left\{ \mathrm{Ext}^1_{\mathrm{U}(L)}(k,k) \otimes \mathrm{Ext}^1_{\mathrm{U}(L)}(k,k) \to (\mathrm{Ext}^1_{\mathrm{U}(L)}(k,k))^2 \right\} = \ker \left\{ H_{(1)} \otimes H_{(1)} \to (H_{(1)})^2 \right\}$$

and hence the 1-2-algebra of $\mathrm{Ext}_{\mathrm{U}(L)}(k,k)$ is equal to the 1-2-algebra of H.

Third proof.

Let (K,d) be the Koszul complex of A with respect to $(x_1, \dots, x_n)$; i.e., $(K,d) = A[sx_1, \dots, sx_n;\ d(sx_i) = x_i]$ (we could also have used $(\bigwedge(V_{\geq 1}), d)$ from above). Then (K,d) is a differential graded algebra (a DGA) and $\mathrm{H}(K,d) = H$. Since k is a field, H is a strong deformation retract of (K,d) (an SDR); i.e., there are maps $f\colon (K,d) \to (H,0)$ and $\nabla\colon (H,0) \to (K,d)$ such that $f\nabla = id$ and ∇f is homotopic to id. From this it is possible to define an SDR $(\mathrm{T}((sK_+)^*), d_K) \to (\mathrm{T}((sH_+)^*), 0)$, where d_K is the tensor extension of the differential on K. Now, the multiplication on K_+ defines a differential d_2 on $\mathrm{T}((sK_+)^*)$ in the same way as in Section 2; i.e., as the composition

$$(sK_+)^* \to K_+^* \to (K_+ \otimes K_+)^* \overset{\phi^{-1}}{\to} (sK_+)^* \otimes (sK_+)^*$$

and $(\mathrm{T}((sK_+)^*), d_K + d_2)$ is the cobar construction on K^*. Thinking of $d_K + d_2$ as a perturbation of d_K, one may use the perturbation lemma to obtain a differential d on $\mathrm{T}((sH_+)^*)$, such that $(\mathrm{T}((sH_+)^*), d)$ is an SDR of $(\mathrm{T}((sK_+)^*), d_K + d_2)$. Moreover d is a derivation (see [Jo-La 00]). But the cohomology of the cobar construction on K^* is $\mathrm{Ext}_{(K,d)}(k,k)$ and this is $\mathrm{U}(\mathfrak{g}_A^{\geq 2})$ ([Av 74]).

Hence we have a DGA, $(\mathrm{T}((sH_+)^*), d)$, with cohomology $\mathrm{U}(\mathfrak{g}_A^{\geq 2})$, which is approximated by $(\mathrm{T}((sH_+)^*), d_2) = \mathrm{Ext}_H(k,k)$. This is the "SDR-version" of Avramov's spectral sequence. (Compare this with the model $(\bigwedge(V_{\geq 1}), d)$ above, which has homology H, approximated by $\mathrm{H}(\bigwedge(V_{\geq 1}), d_2) = \mathrm{Ext}_{\mathrm{U}(\mathfrak{g}_A^{\geq 2})}(k,k)$.)

Put $V^{i,j} = (sH_{i-1,j})^*$ for $i > 1$ and put $V = \oplus V^{i,j}$. Then d is determined by the value of d on V, $d = d_2 + d_3 + \cdots$, where $d_n\colon V \to V^{\otimes n}$, d_2 is (in principal) the dual of the multiplication on H and the higher d_n are duals of Massey products. We have $V^{i,j} = 0$ if $i > j$ and hence the elements of $V^{i,i}$ are cycles. The cohomology of $(\mathrm{T}(V), d)$ in degree (i,i) is $\mathrm{U}(L)^i$ and hence

$$\mathrm{U}(L)^i = V^{i,i} \oplus (V \otimes V)^{i,i}/\text{boundaries} \oplus (V \otimes V \otimes V)^{i,i}/\text{boundaries} \oplus \cdots$$

This decomposition corresponds to the filtration of $\mathrm{U}(L)$ with respect to powers of $I = \mathrm{U}(L)^+$. This is so, because all elements of $(V \otimes V \otimes \cdots \otimes V)^{i,i}/\text{boundaries}$ are products of elements from $V^{j,j}$. In particular, $I/I^2 = \oplus_i V^{i,i} = (sH_{(1)})^*$, $I^2/I^3 = \oplus_i (V \otimes V/d_2(V))^{i,i}$ and

$$\ker \left(I/I^2 \otimes I/I^2 \to I^2/I^3 \right) = \mathrm{im}(d_2) = \phi^{-1} \left(\ker(H_{(1)} \otimes H_{(1)} \to H_{(2)}) \right)^{\perp} .$$

Hence the 1-2-algebra of H is isomorphic to the Koszul dual of $\mathrm{gr}(\mathrm{U}(L))$. Thus, using Proposition 2.4, we get (i) and (ii); (iii) follows as before.

Suppose A is Koszul. Then $(\mathrm{T}(V), d)$ has cohomology only on the diagonal. Consider $j - i$ as the homological degree of $\mathrm{T}(V)^{i,j}$. We have a map $(\mathrm{T}(V), d) \to (\mathrm{U}(L), 0)$ (where $\mathrm{U}(L)$ is considered to be concentrated in homological degree zero), which induces an isomorphism in homology.

Hence $(\mathrm{T}(V), d)$ is a minimal (free associative) model of $\mathrm{U}(L)$. Such a model may always be constructed for any $\mathbb{Z}_2 \times \mathbb{N}$-graded connected k-algebra B. This is similar to the construction of models $(\bigwedge(V), d)$ of commutative algebras described above, but now the space $(sV)^*$ is the algebra $\mathrm{Ext}_B^+(k,k)$ and the dual of d_2 (with the usual sign corrections) is the dual of the Yoneda product. Hence in our case we get $\mathrm{Ext}_{\mathrm{U}(L)}(k,k) \cong H$, which proves (iv). $\square$

Problem 4.5 Is the converse of (iv) in Theorem 4.1 true? Or, perhaps weaker, if the double Poincaré series of $\mathrm{U}(L)$, $P_{\mathrm{U}(L)}(x,y)$, is equal to $H(1/x, xy)$, is it then true that A is Koszul?

Observe that the equality $P_{\mathrm{U}(L)}(x,y) = H(1/x, xy)$ implies the equality $A(z) \cdot A^!(-z) = 1$.

5 The coordinate ring of generic points – IGC and MRC

If $X = \{P_1, \dots, P_s\}$ is a set of points in P_k^n, we denote the coordinate ring of X, $k[X_0, \dots, X_n]/I(X)$, by A_X. It is well-known [Ge-Or 81] that the Hilbert series of A_X satisfies $A_X(z) \leq \sum_{i \geq 0} \min(\binom{n+i}{i}, s) z^i$. There is equality on an open non-empty set in $(P_k^n)^s$, and in case of equality the points are said to be in generic position. A point $P = (c_0 : \cdots : c_n) \in P_k^n$ is called generic if the c_i's are algebraically independent over the prime field of k. A set of mutually generic points is of course in generic position. We will in the sequel always assume that our sets of points are generic. We may also assume that $s \geq n+1$, since otherwise the points lie in P_k^{n-1}.

Definition 5.1 *Given n and $s \geq n+1$, let $d \geq 2$ as a function of n and s be defined by*

$$\binom{n+d-1}{d-1} \leq s < \binom{n+d}{d}.$$

It is well-known that

$$\mathrm{Tor}_{i,j}^S(A_X, k) = 0 \quad \text{if} \quad j \neq i+d-1, i+d$$

(here and in the sequel S will always denote $k[X_0, \dots, X_n]$). In particular the minimal generators for the ideal I_X are of degrees d and $d+1$. Also, it is well-known that if $s = \binom{n+d-1}{d-1}$, then $\mathrm{Tor}_{i,j}^S(A_X, k) = 0$ if $j \neq i+d-1$ (A_X has a linear resolution over S).

It has been conjectured that the map $((I_X)_d)^{n+1} \to (I_X)_{d+1}$, defined by $(f_0, \dots, f_n) \mapsto \sum_{i=0}^n X_i f_i$, is either injective or surjective. This is called the Ideal Generating Conjecture (IGC). It is equivalent to the fact that

$$\mathrm{Tor}_{1,d+1}^S(A_X, k) = 0 \quad \text{or} \quad \mathrm{Tor}_{2,d+1}^S(A_X, k) = 0.$$

The generalization that for all i,

$$\mathrm{Tor}^S_{i,d+i}(A_X,k)=0 \quad \text{or} \quad \mathrm{Tor}^S_{i+1,d+i}(A_X,k)=0$$

is called the Minimal Resolution Conjecture (MRC).

The MRC is proved for P^2 [Ge-Ma 84]; for P^3 [Ba-Ge 86]; for P^4 [Wa 95], [La 97]; for P^n, $n+1\le s\le n+4$ [Ge-Lo 89], [Ca-Ro-Va 91]; for P^n, $s=\binom{n+2}{2}-n$ [Lo 93]; for P^n, $n\le 9$, $s\le 50$ except $(n,s)=(6,11),(7,12),(8,13)$ [Be-Kr 94]; and for P^n, $s>6^{n^3\log n}$ [Hi-Si 96]. There has some time been computational evidence for some counterexamples by Schreyer, later [Bo 94], and [Be-Kr 94], and now there is shown to exist, for each $n\ge 6$, $n\ne 9$, one s which gives a counterexample to MRC [Ei-Po 99]. In [Ge-Gr-Ro 86, Theorems 4.7 and 5.8] IGC is proved in several cases, e.g. for $s\le 1+n+n^2/4$ and for $\binom{n+2}{2}-n\le s<\binom{n+2}{2}$. In [Co-Tr-Va 97] it is proved that A_X is even Koszul if $s\le 1+n+n^2/4$.

We will now state some well-known facts about the Hilbert series of A_X. Let $h^{n,s}_{i,j}=\dim_k \mathrm{Tor}^S_{i,j}(A_X,k)$, $h^{n,s}_{i,j}=h_{i,j}$ when n and s are considered to be fixed and $H(x,y)=\sum h_{i,j}x^iy^j$.

Proposition 5.2 *For any given n and s we have*

$$(i)\qquad A_X(z)=\sum_{j=0}^{d-1}\binom{n+j}{j}z^j+\frac{sz^d}{1-z}$$

$$(ii)\qquad (1-z)^{n+1}A_X(z)=1+\sum_{j=0}^{n}(-1)^{j+1}\binom{n}{j}\left\{\binom{n+d}{d}\frac{d}{d+j}-s\right\}z^{d+j}$$

$$(iii)\qquad (1-z)^{n+1}A_X(z)=H(-1,z)$$

$$(iv)\qquad h_{j+1,d+j}-h_{j,d+j}=\binom{n}{j}\left\{\binom{n+d}{d}\frac{d}{d+j}-s\right\},\quad 0\le j\le n.$$

Proof. The formulas (i) and (iii) are well-known and (iv) follows from (ii) and (iii) and the remark above. To prove (ii) it is enough to prove the identity

$$(1-z)^{n+1}\sum_{j=0}^{d-1}\binom{n+j}{j}z^j=1+\sum_{j=0}^{n}(-1)^{j+1}\binom{n}{j}\binom{n+d}{d}\frac{d}{d+j}z^{d+j}.$$

This can easily be done by proving that both sides have the same derivative. Here is another way to deduce the identity without knowing the right hand side beforehand. Call the left hand side $p(z)$. It is a polynomial of degree $n+d$, divisible by $(1-z)^{n+1}$, and it may also be written as

$$1-(1-z)^{n+1}\sum_{j=d}^{\infty}\binom{n+j}{j}z^j,$$

which shows that $p(z)-1$ is divisible (as a polynomial) by z^d. It follows that $p'(z)$ is divisible by z^{d-1} and by $(1-z)^n$ and hence there is a constant c such that $p'(z)=cz^{d-1}(1-z)^n=c\sum_{j=0}^{n}(-1)^j\binom{n}{j}z^{j+d-1}$. Intergrating this one gets

$$p(z) = 1 + c\sum_{j=0}^{n}(-1)^j \binom{n}{j}\frac{1}{d+j}z^{j+d}\,.$$

But the highest coefficient in $p(z)$ is $(-1)^{n+1}\binom{n+d-1}{d-1}$ and hence $c(-1)^n\frac{1}{d+n} = (-1)^{n+1}\binom{n+d-1}{d-1}$, which gives $c = -d\binom{n+d}{d}$. □

Definition 5.3 *Let n and s be given and d defined by Definition 5.1. Then m, as a function of n and s, is defined as the least integer such that*

$$\binom{n+d}{d}\frac{d}{d+m} - s \le 0$$

and t, as a function of n and s, is defined by

$$t = \binom{n}{m-1}\left\{\binom{n+d}{d}\frac{d}{d+m-1} - s\right\}\,.$$

It is easy to see that if $\mathrm{Tor}^S_{i,d+i-1}(A_X,k) = 0$, we have $\mathrm{Tor}^S_{j,d+j-1}(A_X,k) = 0$ for all $j \ge i$, and if $\mathrm{Tor}^S_{i,d+i}(A_X,k) = 0$, we have $\mathrm{Tor}^S_{j,d+j}(A_X,k) = 0$ for all $j \le i$. From this and Proposition 5.2 the following proposition is easily deduced.

Proposition 5.4 *For any given n and s the following are equivalent.*

(i)	*MRC holds*
(ii)	$h_{j-1,j+d-1} = h_{j+1,j+d} = 0$ *for some* j
(iii)	$h_{m-1,m+d-1} = h_{m+1,m+d} = 0$
(iv)	$h_{j,j+d-1} = \binom{n}{j-1}\left\{\binom{n+d}{d}\frac{d}{d+j-1} - s\right\}$ *and* $h_{j+1,j+d} = 0$ *for some* j
(v)	$h_{m,m+d-1} = t$ *and* $h_{m+1,m+d} = 0$

□

From Proposition 5.2 the following characterization of IGC immediately follows.

Proposition 5.5 *For any given n and s, IGC holds if and only if*

$$h_{2,d+1} = \max(n\left\{\binom{n+d}{d}\frac{d}{d+1} - s\right\}, 0)\,.$$

□

It is possible to interprete MRC and IGC in terms of the homotopy Lie algebra of A_X (if $d > 2$) or the Lie algebra associated to A_X (if $d = 2$).

Theorem 5.6 *Let X be a set of s generic points in P^n_k and let A_X be the coordinate ring of X. Let $\mathfrak{g}$ be the homotopy Lie algebra of A_X and let L be the Lie algebra associated to A_X (see Definition 3.1). Let d, m and t be the*

numbers defined from n *and* s *by Definition 5.1 and 5.3. Then, if* $d > 2$, *we have the following*

$$(i) \qquad \textit{MRC holds} \iff \dim_k \mathfrak{g}^{m+1,m+d-1} = t \quad \textit{and} \quad \mathfrak{g}^{m+2,m+d} = 0$$

$$(ii) \qquad \textit{IGC holds} \iff \dim_k \mathfrak{g}^{3,d+1} = \max(n \left\{ \binom{n+d}{d} \frac{d}{d+1} - s \right\}, 0)$$

If $d = 2$, *we have the following*

$$(iii) \qquad \textit{MRC holds} \iff \dim_k (L/[L,L])^{m+1} = t \quad \textit{and} \quad (L/[L,L])^{m+2} = 0$$

$$(iv) \qquad \textit{IGC holds} \iff \dim_k L^3 = \max(n \left\{ \frac{(n+2)(n+1)}{3} - s \right\}, 0)$$

Proof. Suppose first that $d > 2$. In the third proof of theorem 4.1 it was proved that $\mathrm{U}(\mathfrak{g}_A^{\geq 2})$ is the cohomology of $(\mathrm{T}(V), d)$, where $V^{ij} = (sH_{i-1,j})^*$ for $i > 1$ (and zero otherwise). Moreover the image of the differential is contained in $\oplus_{n \geq 2} V^{\otimes n}$. In our case, we have that $V^{ij} = 0$ if $j - i \neq d-2, d-1$. The differential maps V^{ij} to $\mathrm{T}(V)^{i+1,j}$. Hence, if this is nonzero, $d-2$ or $d-3$ must be a sum of at least two numbers each of which is $d-1$ or $d-2$. This is impossible if $d > 2$. Hence the differential is zero and $\mathrm{U}(\mathfrak{g}_A^{\geq 2}) \cong \mathrm{T}(V)$. In particular $\mathfrak{g}_A^{i+1,i+d-1} \cong (H_{i,i+d-1})^*$ and $\mathfrak{g}_A^{i+1,i+d} \cong (H_{i,i+d})^*$. Now the statements (i) and (ii) follow from Proposition 5.4 (v) and Proposition 5.5.

Suppose that $d = 2$. The statements (iii) and (iv) follow from the same propositions as above and Theorem 4.1 (ii). □

The next theorem shows how the numbers $h_{i,j}$ are changed when the number of points is increased. For a given d, $h_{i,i+d}$ increases with s, while $h_{i+1,i+d}$ decreases. Moreover the rate of change is bounded by $\binom{n}{i}$.

Theorem 5.7 *Let* A_X *be the coordinate ring of a set* X *of* s *generic points in* P_k^n *and let* $h_{i,j}^{n,s} = \dim_k \mathrm{Tor}_{i,j}^S(A_X, k)$. *Then for any* i, *we have*

$$h_{i,i+d}^{n,s'} \leq h_{i,i+d}^{n,s} \leq h_{i,i+d}^{n,s'} + \binom{n}{i}(s - s') \quad \textit{and}$$

$$h_{i+1,i+d}^{n,s} \leq h_{i+1,i+d}^{n,s'} \leq h_{i+1,i+d}^{n,s} + \binom{n}{i}(s - s')$$

whenever s *and* s' *define the same* d; *i.e., whenever* $\binom{n+d-1}{d-1} \leq s' \leq s \leq \binom{n+d}{d}$.

Proof. Let $Y = \{P_1, \ldots, P_s\}$ be a set of s generic points and let $X = Y \backslash \{P_s\}$. After a coordinate change we may assume that $P_s = (1 : 0 : \cdots : 0)$. Let $K = I_X/I_Y$ be the kernel of the natural map $A_Y \to A_X$. We get $K(z) = A_Y(z) - A_X(z) = z^d/(1-z)$, so K_r is one-dimensional in each degree $r \geq d$. Let $f_d + I_Y \in K$, where $f_d \notin I_Y$ is homogeneous of degree d; i.e., $f_d(P_j) = 0$ for $j = 1, \ldots, s-1$ and $f_d(P_s) \neq 0$. Then $(X_i f_d)(P_j) = 0$ for all j and $i = 1, \ldots, n$, and hence $X_i f_d \in I_Y$ for $i = 1, \ldots, n$. Moreover $(X_0^r f_d)(P_j) = 0$ for $j = 1, \ldots, s-1$, and $(X_0^r f_d)(P_s) = f_d(P_s) \neq 0$ for every $r \geq 0$. Thus K_{r+d} is generated by $X_0^r f_d + I_Y$. It follows that K is generated by

f_d as an S-module and that $K \cong S/(X_1,\dots,X_n)[-d]$. We now consider the long exact $\mathrm{Tor}^S_{*,i+d}(*,k)$-sequence of $0 \longrightarrow K \longrightarrow A_Y \longrightarrow A_X \longrightarrow 0$. Since $K \cong S/(X_1,\dots,X_n)[-d]$, we have $\mathrm{Tor}^S_{j,i+d}(K,k) = 0$ if $j \neq i$ and $\mathrm{Tor}^S_{i,i+d}(K,k) = k^{\binom{n}{i}}$, and we get an exact sequence

$$0 \longrightarrow \mathrm{Tor}^S_{i+1,i+d}(A_Y,k) \longrightarrow \mathrm{Tor}^S_{i+1,i+d}(A_X,k) \longrightarrow \mathrm{Tor}^S_{i,i+d}(K,k) \longrightarrow$$

$$\mathrm{Tor}^S_{i,i+d}(A_Y,k) \longrightarrow \mathrm{Tor}^S_{i,i+d}(A_X,k) \longrightarrow 0.$$

The rightmost part of this sequence gives $h^{n,s-1}_{i,i+d} \leq h^{n,s}_{i,i+d} \leq h^{n,s-1}_{i,i+d} + \binom{n}{i}$, and the leftmost part gives $h^{n,s}_{i+1,i+d} \leq h^{n,s-1}_{i+1,i+d} \leq h^{n,s}_{i+1,i+d} + \binom{n}{i}$. The statement now follows by induction. □

Suppose that MRC holds for s' and s'' points in P^n_k. Then, using the previous theorem, we may in some cases conclude that MRC also holds for all s between s' and s''.

Theorem 5.8 *Suppose MRC holds for s' and s'' points in P^n_k, where for some $d \geq 2$ and some j, $1 \leq j \leq n$,*

$$\binom{n+d}{d}\frac{d}{d+j} \leq s' < s'' \leq \binom{n+d}{d}\frac{d}{d+j-1}.$$

Then MRC holds for any s points in P^n_k, where $s' \leq s \leq s''$.

Proof. By Theorem 5.4, we have $h_{j+1,j+d} = 0$ for s' points ($m = j$ for s'). Hence, by Theorem 5.7, $h_{j+1,j+d} = 0$ is true also for s points. Also by Theorem 5.4, we have $h_{j-1,j+d-1} = 0$ for s'' points. This is true if $s'' < \binom{n+d}{d}\frac{d}{d+j-1}$, since then $m = j$ for s''. If $s'' = \binom{n+d}{d}\frac{d}{d+j-1}$, then by Proposition 5.2 for s'' points, $h_{j,d+j-1} - h_{j-1,d+j-1} = 0$ and since MRC holds we must have $h_{j,d+j-1} = h_{j-1,d+j-1} = 0$. Hence, by Theorem 5.7, $h_{j-1,j+d-1} = 0$ is true also for s points. Hence for s points we have $h_{j+1,j+d} = h_{j-1,j+d-1} = 0$, and hence MRC holds for s points. □

Corollary 5.9 *In order to prove MRC for s in the interval $\binom{n+d-1}{d-1} \leq s < \binom{n+d}{d}$, it is enough to prove MRC for $s_j = \lceil \binom{n+d}{d}\frac{d}{d+j} \rceil$, $j = 1,\dots,n-1$, and for $s'_j = \lfloor \binom{n+d}{d}\frac{d}{d+j} \rfloor$. In order to prove IGC in the same interval, it is enough to prove IGC for $s = s_1$ and $s = s'_1$.*

□

Until now the results in this section hold for any d. However, from now on, we will specialize to the case $d = 2$. Using Theorem 5.8 and a result by Anna Lorenzini we can prove MRC in the interval $[\binom{n+2}{2} - n, \binom{n+2}{2})]$ for all n. This also follows from the results in [Ge-Gr-Ro 86].

Corollary 5.10 *If $\binom{n+2}{2} - n \leq s \leq \binom{n+2}{2}$ then MRC holds.*

Proof. MRC is proved in [Lo 93] for $s = \binom{n+2}{2} - n$. Since MRC also holds for $s = \binom{n+2}{2}$ and $\binom{n+2}{2}\frac{2}{3} \leq \binom{n+2}{2} - n \leq \binom{n+2}{2}$ the statement follows from Theorem 5.8. □

The following theorem will be useful in the next section, where we present results of calculations in the Lie algebra associated to A_X.

Theorem 5.11 *Let X be a set of s generic points in P_k^n and let L be the Lie algebra associated to the coordinate ring of X (see Definition 3.1). Suppose $s < \binom{n+2}{2}$. We then have*

(i) $L^3 = 0 \Rightarrow$ *MRC (and hence also IGC) holds for s' points in P_k^n, where $s \leq s' \leq \binom{n+2}{2}$*

(ii) $\dim_k L^3 = n\left\{\frac{(n+2)(n+1)}{3} - s\right\} \Rightarrow$ *IGC holds for s' points in P_k^n, where $s' \leq s$*

(iii) $\dim_k L^3 = n\left\{\frac{(n+2)(n+1)}{3} - s\right\}$ *and* $L^4 = [L^2, L^2] \Rightarrow$ *MRC holds*

Proof. Suppose $L^3 = 0$. By Corollary 4.3 (iii), we get $h_{2,3} = 0$. Since also $h_{0,2} = 0$, MRC holds for s points by Proposition 5.4 (ii). Now MRC also holds for $s'' = \binom{n+2}{2}$ points. Hence, (i) follows from Theorem 5.8.

Suppose $\dim_k L^3 = n\left\{\frac{(n+2)(n+1)}{3} - s\right\}$. Then, by Proposition 5.2 (iv) ($d = 2$ and $j = 1$), we get $h_{2,3} - h_{1,3} = \dim_k L^3$. But $\dim_k L^3 = h_{2,3}$ and hence $h_{1,3} = 0$. By Theorem 5.7, $h_{1,3} = 0$ is true for any s' points in P_k^n, where $n + 1 \leq s' \leq s$ and hence (ii) follows.

The last statement follows, since as above $h_{1,3} = 0$ and, by Corollary 4.3 (iv), $h_{3,4} = 0$. Now use Proposition 5.4 (ii). □

6 Results of calculations

We note that MRC is proved for $s < \binom{n+2}{2}$ if $n < 9$, except for $(n, s) = (6, 11), (7, 12), (8, 13)$, and also for $s \leq 50$ if $n = 9$, [Be-Kr 94]. Thus we treat only the case $n \geq 9$.

Theorem 6.1 *For $n \leq 20$ and $(n+1)(n+2)/3 \leq s < (n+1)(n+2)/2$ MRC holds.*

Proof. Since $s < (n+1)(n+2)/2$ we have $d = 2$. From the table in Section 11 we get that $L^3 = 0$ for $n \leq 20$ and $s = \lceil (n+1)(n+2)/3 \rceil$. Hence the result follows from Theorem 5.11 (i). □

Theorem 6.2 *If $n \leq 20$ and $s < (n+1)(n+2)/2$, IGC holds.*

Proof. If $s \geq (n+2)(n+1)/3$ the conclusion follows from the previous theorem. From the table in Section 11 we get that $\dim_k L^3 = n\left\{\frac{(n+2)(n+1)}{3} - s\right\}$ for $s = \lfloor (n+1)(n+2)/3 \rfloor$. Hence the result follows from Theorem 5.9. □

Definition 6.3

$$
\begin{aligned}
l_3 &= \lceil (n+1)(n+2)/3 \rceil \\
l_4 &= \left\lceil n^2 + n + 1/2 - \frac{1}{2}\sqrt{2n^4 - 2n^2 + 1} \right\rceil \\
l_5 &= \left\lceil n^2/2 + n + 1 - \frac{1}{10}\sqrt{5n^4 + 20} \right\rceil
\end{aligned}
$$

We conjecture that if $s \geq l_i$, then $L^i = 0$, $i = 3, 4, 5$, cf. Conjecture 8.1. From the table in Section 11, it follows that the conjecture is true for L^3 if $n \leq 20$, for L^4, if $n \leq 17$, and for L^5 if $n \leq 11$. We can use the result for L^4 to extend the interval where we can prove MRC.

Theorem 6.4 *If $n \leq 17$ and $l_4 \leq s \leq \binom{n+2}{2}$ then MRC holds.*

Proof. From the table in Section 11 we get that $L^4 = 0$ for $s = l_4$. Hence by Theorem 5.7 it is true also for $s \geq l_4$. Corollary 4.3 gives $h_{3,4} = 0$ for $s \geq l_4$. Since we have proved that $\dim_k L^3 = n\left\{\frac{(n+2)(n+1)}{3} - s\right\}$ for $s = \lfloor (n+1)(n+2)/3 \rfloor$, it follows that $h_{1,3} = 0$ for that many points and then, by Theorem 5.7, it follows that $h_{1,3} = 0$ for $s \leq (n+1)(n+2)/3$. Hence $h_{1,3} = h_{3,4} = 0$ for $l_4 \leq s \leq (n+1)(n+2)/3$ and hence MRC holds in this interval. But by Theorem 6.1, MRC holds in the interval $l_3 \leq s \leq \binom{n+2}{2}$. □

Using Theorem 5.6 (iii), we can show MRC in a few more cases.

Theorem 6.5 *MRC is true if*
a) $n = 9$ and $d = 2$ (i.e., $s < 55$).
b) $(n, s) = (10, 39), (10, 40), (11, 47)$, and $(12, 55)$.

Proof. For $n = 9$ and $s \leq 50$, MRC is proved in [Be-Kr 94]. For $50 < s < 55$ we have proved that $L^3 = 0$ (see Section 11) and hence MRC holds by Theorem 5.11. For the cases in b) we have used the Mathematica program "liedim" to show that $[L^2, L^2] = L^4$. Then, since $\dim_k L^3 = t$, the statement follows from Theorem 5.6. (In fact, the case $(10, 40)$ may be derived from $(10, 39)$ and $(10, 43)$ using Theorem 5.8.) □

It is natural to conjecture that

$$\mathrm{Tor}^S_{r,s}(A_X, k) \wedge \mathrm{Tor}^S_{t,u}(A_X, k) \to \mathrm{Tor}^S_{r+t,s+u}(A_X, k)$$

is either injective or surjective. We can prove this in some special cases.

Theorem 6.6 *Let A_X be the coordinate ring of s generic points in P^n_k and let $S = k[x_0, \ldots, x_n]$. If $n \leq 17$ and $s \geq l_4$ the multiplication*

$$\mathrm{Tor}^S_{1,2}(A_X, k) \wedge \mathrm{Tor}^S_{1,2}(A_X, k) \to \mathrm{Tor}^S_{2,4}(A_X, k)$$

is injective.

Proof. We have shown that $L^4 = 0$ in this interval (see Section 11). Thus Corollary 4.3 (v) gives that the multiplication is injective. □

Theorem 6.7 *Let A_X be the coordinate ring of s generic points in P^n_k and let $S = k[x_0, \ldots, x_n]$. For $(n, s) =$*

$$(3, 6), (5, 13), (6, 17), (6, 18), (7, 21), (7, 22), (8, 26), (8, 27), (9, 32), (9, 33),$$
$$(10, 39), (10, 40), (11, 47), (12, 55)$$

the multiplication

$$\mathrm{Tor}^S_{1,2}(A_X, k) \wedge \mathrm{Tor}^S_{1,2}(A_X, k) \to \mathrm{Tor}^S_{2,4}(A_X, k)$$

is surjective.

Proof. For these pairs of (n,s), we have that $\mathrm{Tor}^S_{i,*}(A_X,k)$ for $i \leq 3$ have the expected dimensions for MRC ([Be-Kr 94] and Theorem 6.5). By Corollary 4.3 (v), the kernel of the multiplication map in the theorem has dimension $\dim_k[L^2,L^2]$, and by Corollary 4.3 (iv), $h_{3,4} = \dim_k L^4/[L^2,L^2]$. Since MRC holds, we know $h_{1,2}$, $h_{2,4}$, and $h_{3,4}$. From Section 11 we know $\dim_k L^4$ for all pairs (n,s) in the theorem. Thus the statement follows by computing dimensions. E.g., if $n=6$ and $s=17$ we have that $h_{1,2} = \dim_k L^2 = 11$ and hence $\dim_k(\mathrm{Tor}^S_{1,2}(A_X,k) \wedge \mathrm{Tor}^S_{1,2}(A_X,k)) = 55$. Also, $h_{3,4} = 0$ since MRC holds and $m=2$, and, by Proposition 5.2 (iv), we get $h_{2,4} = 45$. Hence by Corollary 4.3 (iv), $L^4 = [L^2,L^2]$, and from the table, $\dim_k L^4 = 10$. Hence the map is surjective, since the kernel is 10-dimensional by Corollary 4.3 (v). □

We can check the conjecture about the multiplication in a few more cases. If $s = n+2$, it is easy to see that A_X is Gorenstein. By reasons of degree, the only possible nonsurjective multiplication is $\mathrm{Tor}_i \wedge \mathrm{Tor}_{n-i} \to \mathrm{Tor}_n$. This is surjective, since A_X is Gorenstein.

In P^3 the multiplication is injective or surjective for any choice of degrees and number of points. If $s \geq 7$ this follows from Theorem 6.6. If $s=5$, A_X is Gorenstein. If $s=4$ the resolution is linear and the multiplication trivial. For the remaining case $s=6$, the multiplication

$$\mathrm{Tor}^S_{1,2}(A_X,k) \wedge \mathrm{Tor}^S_{1,2}(A_X,k) \to \mathrm{Tor}^S_{2,4}(A_X,k)$$

is surjective by Theorem 6.7. That the multiplication

$$\mathrm{Tor}^S_{1,2}(A_X,k) \otimes \mathrm{Tor}^S_{2,3}(A_X,k) \to \mathrm{Tor}^S_{3,5}(A_X,k)$$

is surjective follows in a similar way as in the proof of Theorem 6.7 by determining $\dim_k L^5$ (cf. Section 11).

7 Ideals generated by generic quadratic forms

We call a quadratic form $\sum_{1\leq i\leq j\leq n} c_{ij}x_ix_j$ in $k[x_1,\ldots,x_n]$ *generic* if $\{c_{ij}\}$ is algebraically independent over the prime field of k (the variables x_i are considered to be even). We will now consider ideals $(f_1,\ldots,f_t)$ in $k[x_1,\ldots,x_n]$, where $f_k = \sum_{1\leq i\leq j\leq n} c^{(k)}_{ij}x_ix_j$ and $\{c^{(k)}_{ij}\}$ is algebraically independent over the prime field of k. For simplicity we say that such an ideal is generated by t generic quadratic forms. We will determine for which n and t we have that $k[x_1,\ldots,x_n]/I$ is a Koszul algebra if I is generated by t quadratic generic forms. We will identify an ideal $(g_1,\ldots,g_t)$ generated by linearly independent quadratic elements (and thus the quadratic algebra $k[x_1,\ldots,x_n]/(g_1,\ldots,g_t)$) with a point in $\mathrm{Grass}(\binom{n+1}{2},t)$. One part of the following theorem is proved in [Co-Tr-Va 97].

Theorem 7.1 *Let $A = k[x_1,\ldots,x_n]/(g_1,\ldots,g_t)$ be a quadratic algebra, where the variables x_i are considered to be even. If $t \leq n$ there is an open non-empty subset of $\mathrm{Grass}(\binom{n+1}{2},t)$ on which A is Koszul. If $t \geq n^2/4 + n/2$ there is a non-empty countable intersection of open sets in $\mathrm{Grass}(\binom{n+1}{2},t)$ on which A*

is Koszul. If $n < t < n^2/4 + n/2$, *then* A *is not Koszul on an open non-empty subset of* $\mathrm{Grass}(\binom{n+1}{2}, t)$. *In particular, if* $A = k[x_1, \ldots, x_n]/I$, *where* I *is generated by* t *generic quadratic forms, then* A *is Koszul if and only if* $t \leq n$ *or* $t \geq n^2/4 + n/2$.

Proof. If $t \leq n$, then almost all A are complete intersections, and complete intersections are Koszul, cf. [Fr-Lö 91].

Now let $t \geq n^2/4 + n/2$. Consider the rings A with t fixed as points in $G = \mathrm{Grass}(\binom{n+1}{2}, t)$. The subset of G corresponding to rings A with $A_3 = 0$ is open. For such A, the Bar complex of k over A is based on the vector space $A_1 \oplus A_2$, which has the fixed dimension $n + \binom{n+1}{2} - t$, and the differential is expressed by means of the multiplication in A. Hence, it is an open condition that the homology is zero in a certain bidegree (i, j), and hence $A_3 = 0$ and $\mathrm{Tor}^A_{i,j}(k, k) = 0$ is an open condition for fixed (i, j). That A is Koszul, i.e., $\mathrm{Tor}^A_{i,j}(k, k) = 0$ if $i \neq j$, is therefore true on an intersection of open sets. It suffices to show that this intersection is non-empty. If $A = k[x_1, \ldots, x_a]/(x_1, \ldots, x_a)^2 \otimes k[x_1, \ldots, x_b]/(x_1, \ldots, x_b)^2$, with $a = \lceil n/2 \rceil$ and $b = \lfloor n/2 \rfloor$, we have an example of a Koszul algebra in case $t = \lceil n^2/4 + n/2 \rceil$, [Fr 75]. If t is larger we could just divide by some more quadratic monomials.

Now let $n < t \leq \binom{n+2}{3}/n$. Then, by [Ho-La 87], there is a non-empty subset of G with $\{x_i g_j\}$ linearly independent, where $I = (g_1, \ldots, g_t)$. By Corollary 4.3 (iii), this gives $L^3 = 0$. If A were Koszul, then $U(\eta_A)$ is equal to $\mathrm{Ext_A}(k, k)$. Then A would be a complete intersection by [Gu 71], a contradiction since $t > n$.

Finally let $\binom{n+2}{3}/n \leq t < n^2/4 + n/2$. Then, by [Ho-La 87], $A_3 = 0$ on a non-empty open subset of G. If A is Koszul, then $A^!(z) = 1/(1 - nz + (\binom{n+1}{2} - t)z^2) = 1/((1 - nz/2)^2 + (n^2/4 + n/2 - t)z^2)$. Hence $A^!(z)$ has no pole on the real axis. But then the coefficients in $A^!(z)$ are not all non-negative, since a series with non-negative coefficients has a singularity on the positive real axis if its radius of convergence is finite. □

Remark 7.2 It follows from the theorem above that the "smallest" algebra defined by generic quadratic forms which is not Koszul is obtained by choosing $n = 4$ and $t = 5$. This was in fact the first example of a non-Koszul quadratic algebra found by Christer Lech 1975.

To state some conjectures about the coordinate ring of generic quadratic forms, we need a concept from non-commutative graded algebras.

Let B be an $\mathbb{N}$-graded associative algebra and $\{y_1, \ldots, y_r\}$ a set of homogeneous elements in B of degrees $d_1, \ldots, d_r$, respectively. Then

$$B/(y_1, \ldots, y_r)(z) \geq B(z)/(1 + (z^{d_1} + \cdots + z^{d_r})B(z))$$

and the set is called *strongly free* in B if there is equality, [An 82]. Any subset of a strongly free set is strongly free. If $\{y_1, \ldots, y_r\}$ is a strongly free set of quadratic elements in the free associative algebra $k\langle X_1, \ldots, X_n\rangle$, we get

$$k\langle X_1, \ldots, X_n\rangle/(y_1, \ldots, y_r)(z) = 1/(1 - nz + rz^2).$$

Such an algebra will be called a non-commutative c.i. (complete intersection). They play much the same rôle for non-commutative algebras as complete intersections do for commutative algebras. E.g., the algebra
$B = k\langle X_1, \dots, X_n\rangle/(y_1, \dots, y_r)$ is a non-commutative c.i. if and only if $k\langle X_1, \dots, X_n, Y_1, \dots Y_r;\ dX_i = 0, dY_i = y_i\rangle$ is a minimal model of B (cf. end of proof of Theorem 4.1). It follows that B is a non-commutative c.i. if and only if $\mathrm{gldim}(B) \leq 2$, and hence a non-commutative c.i. is Koszul. Also, for a quadratic algebra A, $A^!$ is a non-commutative c.i. if and only if A is Koszul and $A_3 = 0$.

We will now consider ideals $(h_1, \dots, h_r)$ in $k\langle X_1, \dots, X_n\rangle$, where the variables X_i are odd and where $h_k = \sum_{1\leq i\leq j\leq n} d_{ij}^{(k)}[X_i, X_j]$ and $[X_i, X_j] = X_iX_j + X_jX_i$. Such an element is called a Hopf element and the corresponding quotient algebra a quadratic Hopf algebra. We will identify an ideal $(h_1, \dots, h_r)$ generated by linearly independent quadratic Hopf elements (and thus also the quadratic Hopf algebra $k\langle X_1, \dots, X_n\rangle/(h_1, \dots, h_r)$) with a point in $\mathrm{Grass}(\binom{n+1}{2}, r)$. As before we will say that the ideal $(h_1, \dots, h_r)$ is generated by r generic quadratic Hopf elements if the coefficients $\{d_{ij}^{(k)}\}$ is algebraically independent over the prime field of k.

There is a well-known duality between Grassmanians yielding the following proposition.

Proposition 7.3 *The correspondence $A \leftrightarrow A^!$ defines an isomorphism of projective manifolds between $\mathrm{Grass}(\binom{n+1}{2}, t)$ and $\mathrm{Grass}(\binom{n+1}{2}, \binom{n+1}{2} - t)$*

Proof. The relations in $A^!$ are obtained from those in A by solving a system of linear equations. By Cramer's rule the solutions may be expressed as rational functions and locally the same determinant may be used in the denominators. □

Now the dual version of Theorem 7.1 directly follows.

Theorem 7.4 *Let $B = k\langle X_1, \dots, X_n\rangle/(h_1, \dots, h_r)$ be a quadratic Hopf algebra, where the variables X_i are considered to be odd. If $r \leq n^2/4$ there is a countable intersection of open sets in $\mathrm{Grass}(\binom{n+1}{2}, r)$ on which B is Koszul. If $r \geq \binom{n}{2}$ there is an open non-empty subset of $\mathrm{Grass}(\binom{n+1}{2}, r)$ on which B is Koszul. If $n^2/4 < r < \binom{n}{2}$, then A is not Koszul on an open non-empty subset of $\mathrm{Grass}(\binom{n+1}{2}, r)$. In particular, if $B = k\langle X_1, \dots, X_n\rangle/J$, where J is generated by r generic quadratic Hopf elements, then B is Koszul if and only if $r \leq n^2/4$ or $r \geq \binom{n}{2}$ and B is a non-commutative c.i. if and only if $r \leq n^2/4$.*

Proof. To get the statement about the Koszul property, we just use Theorem 7.1 and Proposition 7.3, observing that $\binom{n+1}{2} - n^2/4 - n/2 = n^2/4$ and $\binom{n+1}{2} - n = \binom{n}{2}$. The last statement (proved in [An 82]) follows from the remark above that B is a non-commutative c.i. if and only if $B^!$ is Koszul and $B^!_3 = 0$, which by the proof of Theorem 7.1 is true iff $\binom{n+1}{2} - r \geq (n+2)(n+1)/6$ and also $\binom{n+1}{2} - r \geq n^2/4 + n/2$ or $\binom{n+1}{2} - r \leq n$. But for $n \geq 2$ we have $n \leq (n+2)(n+1)/6 \leq n^2/4 + n/2$. □

Conjecture 7.5 *Let* $A = k[x_1, \dots, x_n]/I$, *where the variables* x_i *are even and where* I *is generated by quadratic generic elements. Then either* $A^!$ *is a non-commutative c.i. or* η_A *is nilpotent, where* $A^! = U(\eta_A)$.

For a formal power series $\sum_{i\geq 0} a_i z^i$ we let $(\sum_{i\geq 0} a_i z^i)_+ = \sum_{i\geq 0} b_i z^i$, where $b_i = a_i$ if $a_j > 0$ for all $j \leq i$, and $b_i = 0$ otherwise. A formal power series $p(z)$ with integer coefficients and constant term 1 may be written in the form $p(z) = \prod_{k=1}^{\infty}(1+z^{2k-1})^{e_{2k-1}}/(1-z^{2k})^{e_{2k}}$. We denote by $\mathrm{Log}(p(z))$ the series $\sum_{i\geq 1} e_i z^i$. Recall that, for a graded algebra A, we have $U(\eta_A) = A^!$, so $\eta_A(z) = \mathrm{Log}(A^!(z))$.

We now make the conjecture more precise.

Conjecture 7.6 *Let* $A = k[x_1, \dots, x_n]/I$, *where the variables* x_i *are even and where* I *is generated by* $\binom{n+1}{2} - r$ *quadratic generic forms. Then* $\eta_A(z) = (\mathrm{Log}(1/(1-nz+rz^2)))_+$.

Theorem 7.7 *Conjecture 7.6 is true in the following cases*
a) If A *is Koszul.*
b) If $n \leq 8$.
c) If $r \geq (n^2-1)/3$ *and* $n \leq 20$.
d) If $r \geq n^2 - 1/2 - \frac{1}{2}\sqrt{2n^4-2n^2+1}$ *and* $n \leq 17$.
e) If $r \geq n^2/2 - \frac{1}{10}\sqrt{5n^4+20}$ *and* $n \leq 10$.
f) If $(n,r) = (9,22)$.

Proof. a) According to Theorem 7.1, A is Koszul if and only if $r \geq \binom{n}{2}$ or $r \leq n^2/4$. In the first case we know that A is a complete intersection, so $\eta_A(z) = nz + (\binom{n+1}{2} - r)z^2$. A simple calculation shows that $(\mathrm{Log}(1/(1-nz+rz^2)))_+ = nz + (\binom{n+1}{2} - r)z^2$ if and only if $r \geq (n^2-1)/3$. Since $\binom{n}{2} \geq (n^2-1)/3$, the result follows in the first case. In the second case, $A^!$ is a non-commutative c.i. (since A is Koszul and $A_3 = 0$) and hence $A^!(z) = 1/(1-nz+rz^2)$. It follows that $\eta_A(z) = \mathrm{Log}(A^!(z)) = \mathrm{Log}(1/(1-nz+rz^2))$. Hence also $\eta_A(z) = (\mathrm{Log}(1/(1-nz+rz^2)))_+$, since $(\eta_A)_i = 0$ for some i implies that $(\eta_A)_j = 0$ for all $j \geq i$.

b) We show how the result follows for $n = 8$. Then A is Koszul if $\binom{9}{2} - r = t \geq 8^2/4 + 8/2 = 20$; i.e., if A has at least 20 quadratic relations. If A has t quadratic relations, then $A^!$ has $r = \binom{9}{2} - t = 36 - t$ quadratic relations, so $A^!(z) \geq 1/(1-8z+rz^2)$ and hence $\eta_A(z) \geq (\mathrm{Log}(1/(1-8z+rz^2)))_+$. To see that there is equality generically when $t < 20$, it is sufficient to produce one example. Let $\overline{A} = A_X/y$, where A_X is the coordinate ring of s points in P^8, and y is a linear nonzerodivisor in A_X. Then A_X and A have $\binom{10}{2} - s = 45 - s$ quadratic relations. Thus we can use our calculations for 26 or more points in P^8. For 26 to 30 points we verify the conjecture directly from the table in Section 11. Now $s \geq 30$ is equivalent to $t \leq 15$ and to $r \geq 21$. If $n = 8$ we get $(n^2-1)/3 = 21$, so $(\mathrm{Log}(1/(1-nz+rz^2))_+ = nz + (\binom{n+1}{2} - r)z^2$, cf. the proof of a). But from the table we get that $(\eta_{\bar{A}})_3 = 0$ for $s = 30$ and hence this is true also for $s > 30$. Hence $\eta_{\bar{A}} = nz + (\binom{n+1}{2} - r)z^2 = (\mathrm{Log}(1/(1-nz+rz^2))_+$ and we have proved the conjecture for all t when $n = 8$. If $n < 8$ the proof is similar.

c) We shall prove the statement for $t \leq \binom{n+1}{2} - (n^2-1)/3 = (n^2+3n+2)/6$ relations. As before it suffices to find one example. We get such an example from

s points in P^n if $\binom{n+2}{2} - s \leq (n^2+3n+2)/6$; i.e., if $s \geq (n^2+3n+2)/3$. We refer to Theorem 8.2 a). Observe that $r = s-n-1$.

d) We shall prove the statement for $t \leq \binom{n+1}{2} - (n^2 - 1/2 - \frac{1}{2}\sqrt{2n^4-2n^2+1}) = (n+1-n^2)/2 + \frac{1}{2}\sqrt{2n^4-2n^2+1}$ relations. As before it suffices to find one example. We get such an example from s points in P^n if $\binom{n+2}{2} - s \leq (n+1-n^2)/2 + \sqrt{2n^4-2n^2+1}$; i.e., if $s \geq n^2+n+\frac{1}{2}+\frac{1}{2}\sqrt{2n^4-2n^2+1}$. We refer to Theorem 8.2 b).

e) As in d) it suffices to consider $s \geq \frac{n^2}{2} + n + 1 - \frac{1}{10}\sqrt{5n^4+20}$ points in P^n. We refer to Theorem 8.2 c).

f) Here we can refer to 8.2 d); i.e., 32 points in P^9. □

We can also give some partial results.

Theorem 7.8 *If A is as in Conjecture 7.6, we have*
a) $\eta_A(z) \equiv (\mathrm{Log}(1/(1-nz+rz^2)))_+ \pmod{z^6}$ *if* $n \leq 10$.
b) $\eta_A(z) \equiv (\mathrm{Log}(1/(1-nz+rz^2)))_+ \pmod{z^5}$ *if* $n \leq 17$.
c) $\eta_A(z) \equiv (\mathrm{Log}(1/(1-nz+rz^2)))_+ \pmod{z^4}$ *if* $n \leq 20$.
d) $\eta_A(z) \equiv (\mathrm{Log}(1/(1-nz+rz^2)))_+ \pmod{z^3}$.

Proof. We use the same method as in the preceeding proof. □

We end this section by reformulating the conjectures above in terms of Hopf algebras.

Conjecture 7.9 *Let $B = k\langle X_1, \ldots, X_n\rangle/J$, where the variables X_i are odd and where J is generated by quadratic generic Hopf elements. Then either B is a non-commutative c.i. or η is nilpotent, where $B = U(\eta)$.*

Conjecture 7.10 *Let $B = k\langle X_1, \ldots, X_n\rangle/J$, where the variables X_i are odd and where J is generated by r quadratic generic Hopf elements. Then $\eta(z) = (\mathrm{Log}(1/(1-nz+rz^2)))_+$, where $B = U(\eta)$.*

Using Proposition 7.3 it is easy to see that these new conjectures are equivalent to their commutative versions above. In fact, it is obvious that the old conjectures imply the new ones, since if B is a quadratic Hopf algebra which is a non-commutative c.i., or its underlying Lie algebra is nilpotent, or it has the minimal possible Hilbert series, then the same is true for the generic case.

Suppose that Conjecture 7.9 is true and consider quadratic commutative algebras A defined by t relations. Since $A^!$ is a non-commutative c.i. if and only if A is Koszul and $A_3 = 0$, it follows from the proof of Theorem 7.1, that the set on which $A^!$ is a non-commutative c.i. is an intersection of countably many open sets in $\mathrm{Grass}(\binom{n+1}{2}, t)$. Also (using Proposition 7.3), the set on which η_A is nilpotent is an open set in $\mathrm{Grass}(\binom{n+1}{2}, t)$. One of these sets must be non-empty, since if B is the generic Hopf algebra defined by $\binom{n+1}{2} - t$ quadratic Hopf elements, then $B^!$ is a member of one of the sets by Conjecture 7.9. But then also the quadratic commutative algebra defined by t generic relations is a member of the corresponding set.

Suppose now that Conjecture 7.10 is true. Then for each r, there is a quadratic commutative algebra A defined by $\binom{n+1}{2} - r$ relations, such that $\eta_A(z)$ is the correct series. But since this is the minimal possible series, it follows from Proposition 7.3 that the set of A in $\mathrm{Grass}(\binom{n+1}{2}, \binom{n+1}{2} - r)$ such that $\eta_A(z)$ has the right value is an intersection of open sets. Since this set is non-empty it contains the generic algebra A.

8 Generic points again

Now let us consider the coordinate ring A of s generic points in P_k^n. Let $\overline{A} = A/y$ where y is a linear nonzerodivisor. If $n+1 \leq s < \binom{n+2}{2}$ then $\overline{A}(z) = 1 + nz + (s-n-1)z^2$. This is the series you get if you divide a polynomial ring in n variables with $\binom{n+2}{2} - s$ generic quadratic forms, where $s \leq (n+2)(n+1)/3$ (see the proof of Theorem 7.1). Let $A^! = \mathrm{U}(\eta_A)$ and $\overline{A}^! = \mathrm{U}(\eta_{\bar{A}})$. It is natural to conjecture that for any $n+1 \leq s < \binom{n+2}{2}$, $\eta_{\bar{A}}$ behaves as the free Lie algebra on n generators divided by $s-n-1$ generic quadratic Lie elements. Hence by Conjecture 7.10, we should have that $\eta_{\bar{A}}(z) = (\mathrm{Log}(1/(1-nz+(s-n-1)z^2)))_+$. But since $A(z) = (1+z)\overline{A}^!(z)$ we should have $\eta_A(z) = (\mathrm{Log}((1+z)/(1-nz+(s-n-1)z^2)))_+$. Hence it is natural to make the following conjecture.

Conjecture 8.1 *Let A be the coordinate ring for s generic points in P_k^n, where $n+1 \leq s < \binom{n+2}{2}$. Let $A^! = \mathrm{U}(\eta_A)$ and $L = \eta_A^{\geq 2}$. Then*

$$\eta_A(z) = (n+1)z + L(z) = (\mathrm{Log}((1+z)/(1-nz+(s-n-1)z^2)))_+.$$

In particular
$\dim_k L^3 = \max((n^3-n)/3 - (s-n-1)n, 0)$.
$\dim_k L^4 = \max((n^4-n^2)/4 - (s-n-1)n^2 + \binom{s-n}{2}, 0)$.
$\dim_k L^5 = \max(n^5/5 + n^4 + 2n^3 + 2n^2 + 4n/5 - s(n^3+2n^2+2n) + ns^2, 0)$.
Hence we conjecture that
$L^3 = 0$ *if and only if* $s \geq l_3$.
$L^4 = 0$ *if and only if* $s \geq l_4$.
$L^5 = 0$ *if and only if* $s \geq l_5$.
(For the definition of l_3, l_4 and l_5, see Definition 6.3.)

Theorem 8.2 *Conjecture 8.1 is true in the following cases:*
a) If $l_3 \leq s < \binom{n+2}{2}$ *and* $n \leq 20$.
b) If $l_4 \leq s < \binom{n+2}{2}$ *and* $n \leq 17$.
c) If $l_5 \leq s < \binom{n+2}{2}$ *and* $n \leq 10$.
d) If $s \leq 1 + n + n^2/4$.
e) If $n \leq 8$.
f) If $(n,s) = (9,32)$.

Remark. The LHS in b) is approximately $0.293n^2$ and the LHS in c) is approximately $0.276n^2$ for large n.

To prove the theorem we need a lemma.

Lemma 8.3 *If $L^i = 0$ for some set of s points in P^n, then $L^i = 0$ for t generic points if $t \geq s$.*

Proof. If s squares generate $\mathcal{L}_i$, then also t generic squares do, if $t \geq s$. □

We need a variation of the concept of strongly free sets. We define a set of quadratic elements $\{y_1, \ldots, y_t\}$ in $k\langle X_1, \ldots, X_n\rangle$ to be d-strongly free if

$$k\langle X_1, \ldots, X_n\rangle/(y_1, \ldots, y_t)(z) \equiv 1/(1-nz+tz^2) \pmod{z^{d+1}}.$$

Lemma 8.4 *Any subset of a d-strongly free set is d-strongly free.*

Proof. This follows from the corresponding proof in [An 82]. □

Proof of Theorem 8.2. We have shown that $L^3 = 0$ for $s = l_3$ random points if $n \leq 20$ (see the table in Section 11). Then Lemma 8.3 gives a). For $t = \lfloor (n+1)(n+2)/3 \rfloor$ random points we have shown that $\dim_k L^3 = -nt + 2\binom{n+2}{3}$ if $n \leq 20$ (see the table). This gives that the t random squares generating $I_X^\perp$ is a 3-strongly free set. Then of course t truly generic squares constitute a 3-strongly free set, and Lemma 8.4 shows that $s \leq t$ generic squares are 3-strongly free, hence $\eta(z) \equiv (\mathrm{Log}((1+z)/(1-nz+(s-n-1)z^2)))_+ \pmod{z^4}$ if $n \leq 20$. We have also shown that $L^4 = 0$ if $s = \lceil l_4 \rceil$ if $n \leq 17$. Hence $L^4 = 0$ if $s \geq l_4$ and $n \leq 17$ by Lemma 8.3. Then $L(z) \equiv (\mathrm{Log}((1+z)/(1-nz+(s-n-1)z^2)))_+ \pmod{z^5}$ if $n \leq 17$ and b) is proved. c) is proved similarly. In [Co-Tr-Va 97] it is proved that A_X is Koszul if $s \leq 1 + n + n^2/4$. This gives that $(A_X)^!(z) = 1/(1-nz+(s-n-1)z^2)$, so d) follows. For e) and f) see the table. □

9 Poincaré series of generic points

In this section we study the Poincaré series of the coordinate ring A_X; that is, $P_{A_X}(x,y) = \sum \dim_k \mathrm{Tor}_{i,j}^{A_X}(k,k) x^i y^j$. Here, we allow any s, not only $s < \binom{n+2}{2}$. In fact, the case $s \geq \binom{n+2}{2}$ will be the easy one.

Conjecture 9.1 *The Poincaré series $P_{A_X}(x,y)$ is rational, where A_X is the coordinate ring of a set X of s generic points in P_k^n.*

Theorem 9.2 *Conjecture 9.1 is true in the following cases.*
a) If $s \geq \binom{n+2}{2}$.
b) If Conjecture 8.1 is true.

Proof. In case a) we may argue as in the proof of Theorem 5.6 to conclude that $P_{A_X}(x,y) = (1-x)^{n+1}/(1-x(H(x,y)-1))$, where $H(x,y)$ is the series of the homology of the Koszul complex (A_X is Golod).

In case b) we use the fact that, if $A = A_X/f$ where f is some linear nonzerodivisor, then $P_A(x,y)$ is rational if and only if $\mathrm{U}(L)(x,y)$ is rational, where L is the Lie algebra associated to A, [Lö 86]. If Conjecture 8.1 is true, then either L is nilpotent (and hence $\mathrm{U}(L)(z)$ is rational) or $\mathrm{U}(L)(z) = \frac{1}{(1+z)^n(1-nz+(s-n-1)z^2)}$ □

10 Generic forms in the exterior algebra

If $f_1, \ldots, f_t$ are generic quadratic forms in $k[x_1, \ldots, x_n]$, k a field of characteristic 0, there is a well-known conjecture for the Hilbert series of $A_{n,t} = k[x_1, \ldots, x_n]/(f_1, \ldots, f_t)$, namely that $A_{n,t}(z) = ((1-z^2)^t/(1-z)^n)_+$. This conjecture is proved in many cases, and no counterexamples are known. When calculating examples of ideals generated by quadratic forms with the expected generic Hilbert series, experience seems to indicate that one does not have to

choose all forms at random. In fact, it seems that one can almost always choose the first n forms to be the squares of the variables. In fact, we know of only one exception, namely seven forms in five variables over a field of characteristic 2. This made us look at the problem of Hilbert series of generic forms in the exterior algebra. It turns out that the answer to the question in the exterior algebra, which is analogous to the Hochster-Laksov result for polynomial rings, is not what we expected. Although $\dim_k \wedge^3(V) = 10$ if V is 5-dimensional, two generic forms of degree 2 do not generate $\wedge^3(V)$.

Theorem 10.1 *Consider the exterior algebra E over a vector space with basis $\{e_1, \dots, e_5\}$ (over any field). Let $f_1 = c_1 e_1 \wedge e_2 + c_2 e_1 \wedge e_3 + c_3 e_1 \wedge e_4 + c_4 e_1 \wedge e_5 + c_5 e_2 \wedge e_3 + c_6 e_2 \wedge e_4 + c_7 e_2 \wedge e_5 + c_8 e_3 \wedge e_4 + c_9 e_3 \wedge e_5 + c_{10} e_4 \wedge e_5$ and $f_2 = c_{11} e_1 \wedge e_2 + c_{12} e_1 \wedge e_1 + c_{13} e_1 \wedge e_4 + c_{14} e_1 \wedge e_5 + c_{15} e_2 \wedge e_3 + c_{16} e_2 \wedge e_4 + c_{17} e_2 \wedge e_5 + c_{18} e_3 \wedge e_4 + c_{19} e_3 \wedge e_5 + c_{20} e_4 \wedge e_5$ be two forms in E. Then $\{e_1 \wedge f_1, \dots, e_5 \wedge f_1, e_1 \wedge f_2, \dots, e_5 \wedge f_2\}$ is linearly dependent.*

Proof. It suffices to prove the theorem for generic forms (so we can suppose that the c_i's are algebraically independent over the prime field of k). In fact,

$(-c_3c_1c_{12}c_{20} - c_3c_1c_{18}c_{14} + c_3c_1c_{19}c_{13} + c_{16}c_3c_2c_{14} - c_{16}c_{12}c_4 + c_3c_2c_{11}c_{20} + c_{12}c_3c_{17} + c_{18}c_4c_{11} - c_{19}c_3c_{11} + c_{13}c_4c_{15} - c_{13}c_3c_2c_{17} - c_3c_{15}c_{14}$,

$c_3c_1c_{16}c_{19} - c_3c_1c_{15}c_{20} - c_3c_1c_{18}c_{17} - c_{16}c_7c_{12} + c_{16}c_5c_{14} + c_5c_{11}c_{20} + c_6c_{17}c_{12} + c_{18}c_7c_{11} + c_{13}c_{15}c_7 - c_5c_{13}c_{17} - c_{15}c_6c_{14} - c_6c_{11}c_{19}$,

$-c_9c_{12}c_{16} + c_3c_2c_{19}c_{16} + c_9c_{11}c_{18} + c_9c_{13}c_{15} + c_{20}c_{12}c_5 - c_{20}c_3c_2c_{15} + c_{12}c_8c_{17} - c_{11}c_8c_{19} - c_3c_2c_{18}c_{17} + c_{18}c_5c_{14} - c_5c_{19}c_{13} - c_8c_{14}c_{15}$,

$-c_{16}c_{12}c_{10} + c_{16}c_3c_{19} - c_8c_{14}c_{16} + c_{12}c_6c_{20} + c_6c_{14}c_{18} + c_8c_{17}c_{13} - c_3c_{20}c_{15} + c_{18}c_{11}c_{10} - c_8c_{11}c_{20} - c_{13}c_6c_{19} + c_{13}c_{15}c_{10} - c_3c_{18}c_{17}$,

$-c_{16}c_{14}c_9 + c_{16}c_4c_{19} - c_9c_{11}c_{20} + c_9c_{17}c_{13} + c_{20}c_7c_{12} - c_{20}c_4c_{15} - c_{10}c_{17}c_{12} + c_{11}c_{10}c_{19} - c_7c_{19}c_{13} - c_4c_{18}c_{17} + c_7c_{18}c_{14} + c_{14}c_{15}c_{10}$,

$-c_3c_1c_{13}c_9 + c_3c_1c_{12}c_{10} + c_3c_1c_8c_{14} + c_3c_9c_{11} + c_6c_4c_{12} - c_{12}c_3c_7 - c_3c_2c_{10}c_{11} - c_8c_{11}c_4 - c_5c_{13}c_4 + c_{13}c_3c_2c_7 - c_6c_3c_2c_{14} + c_3c_5c_{14}$,

$-c_3c_1c_9c_{16} + c_3c_1c_{15}c_{10} + c_3c_1c_8c_{17} - c_{16}c_5c_4 + c_3c_2c_7c_{16} + c_9c_6c_{11} - c_5c_{11}c_{10} + c_6c_{15}c_4 - c_8c_7c_{11} - c_3c_2c_6c_{17} - c_{15}c_7c_3 + c_3c_5c_{17}$,

$-c_3c_1c_{18}c_9 + c_3c_1c_8c_{19} + c_9c_{12}c_6 - c_3c_9c_{15} - c_{12}c_{10}c_5 - c_8c_{12}c_7 - c_5c_{18}c_4 + c_8c_{15}c_4 + c_3c_2c_{18}c_7 - c_6c_3c_2c_{19} + c_5c_3c_{19} + c_3c_2c_{10}c_{15}$,

$c_3c_1c_{20}c_8 - c_3c_1c_{18}c_{10} - c_3c_9c_{16} + c_{16}c_8c_4 + c_3c_2c_{10}c_{16} + c_{13}c_9c_6 + c_5c_{20}c_3 - c_3c_2c_6c_{20} - c_{18}c_6c_4 - c_{13}c_5c_{10} - c_8c_7c_{13} + c_{18}c_7c_3$,

$c_3c_1c_9c_{20} - c_3c_1c_{10}c_{19} - c_9c_3c_{17} + c_9c_{14}c_6 - c_7c_3c_2c_{20} + c_5c_4c_{20} + c_4c_8c_{17} - c_5c_{10}c_{14} - c_{14}c_8c_7 + c_3c_2c_{17}c_{10} + c_7c_3c_{19} - c_4c_6c_{19})$

is a relation. □

Notice that this also gives a proof of the fact that the ideal $(x_1^2, \dots, x_5^2)$ in $S = k[x_1, \dots, x_5]$ can not be extended in any way with two quadratic forms to an ideal I with the expected generic Hilbert series $S/I(z) = 1 + 5z + 8z^2$, if $\mathrm{char}(k) = 2$. We believe, however, that the example above is the only counterexample to the following conjecture.

Conjecture 10.2 *If A is an exterior algebra of a vector space of dimension n modulo an ideal generated by t generic quadratic forms, then $A(z) = ((1+z)^n(1-z^2)^t)_+$.*

If A is an exterior algebra modulo an ideal generated by quadratic forms, then $A^! = U(\eta_A)$, where η_A is an ordinary Lie algebra; i.e., all generators are even. If the defining ideal of A is generated by $\binom{n}{2} - r$ quadratic forms, then η_A is a free Lie algebra on n generators modulo an ideal generated by r quadratic elements. Recall, that if $U(\eta_A) = A^!$, then $\eta_A(z) = \log(A^!(z))$, where $\log(A^!(z)) = \sum_{i\geq 1} e_i z^i$ if $A^!(z) = \prod_{i=1}^{\infty} 1/((1-z^i)^{e_i}$.

One could ask when a quotient of an exterior algebra on n generators modulo t generic quadratic forms is Koszul. A natural guess, based on the results in Section 7, is that this is true if and only if $t \geq \binom{n}{2} - n^2/4$ or $t = 0$. We can prove this in one direction in the same way as in the proof of Theorem 7.1. The other direction is harder, since the example above shows that the Hochster-Laksov result cannot directly be translated to the exterior algebra case.

Theorem 10.3 *Consider an exterior algebra A on n generators modulo t linearly independent quadratic forms as a point in* $\mathrm{Grass}(\binom{n}{2}, t)$. *If $t \geq \binom{n}{2} - n^2/4$ there is a non-empty countable intersection of open sets in* $\mathrm{Grass}(\binom{n}{2}, t)$ *on which A is Koszul. Moreover, A is not Koszul if $t > 0$ and η_A is nilpotent, where $A^! = \mathrm{U}(\eta_A)$.*

Proof. The same argument as in the proof of Theorem 7.1 shows that the set of A in $\mathrm{Grass}(\binom{n}{2}, t)$, such that $A_3 = 0$ and A is Koszul, is a countable intersection of open sets. We may use the same example as before to show that this set is non-empty if $t \geq \binom{n}{2} - n^2/4$. Indeed, if $a = \lceil n/2 \rceil$ and $b = \lfloor n/2 \rfloor$, a simple computation shows that $\binom{a}{2} + \binom{b}{2} = \lceil \binom{n}{2} - n^2/4 \rceil$. If η_A is nilpotent and $(\eta_A)_2 \neq 0$, then $1/A^!(z)$ is a polynomial of degree $> n$ (see above), which is impossible if A is Koszul, since $A(z)$ is a polynomial of degree $\leq n$. □

In analogy to the case where the variables are even, it is natural to make the following conjecture.

Conjecture 10.4 *If A is a generic quadratic quotient of a polynomial algebra, which is commutative in the graded sense, then $A^!$ is either a non-commutative c.i. or η_A is nilpotent.*

The conjecture is false for 3 generic forms in 5 odd variables, see below. We also make the conjecture more precise in the odd case.

Conjecture 10.5 *Let A be an exterior algebra on n variables, modulo $\binom{n}{2} - r$ generic quadratic forms. Then $\eta_A(z) = (\log(1/(1 - nz + rz^2)))_+$.*

If $(n, r) = (5, 7)$ or $(5, 8)$, the conjecture is false, see below. We believe, however, that these are the only counterexamples.

Theorem 10.6 *Conjecture 10.5 is true in the odd case for $n \leq 8$ except for $(n, r) = (5, 7), (5, 8)$ when it is false. It is true also for $n = 9, r \neq 21$ and in general for $r \leq n^2/4$.*

Proof. The case $r \leq n^2/4$ follows from Theorem 10.3, since then $A^!(z) = 1/(1 - nz + rz^2)$. For the other cases, it is only necessary to provide an example with the correct series in each case. We refer to the last table in Section 11. Observe that the conjecture is trivially true when $(\eta_A)^3 = 0$. □

As in Section 7 we may reformulate the two conjectures above in terms of Hopf algebras. Consider the free associative algebra $k\langle X_1, \dots, X_n\rangle$, where the variables have some parities $|X_i| \in \mathbb{Z}_2$. We will call an element $f = \sum c_{ij}[X_i, X_j]$ an Hopf element, where $[X_i, X_j] = X_iX_j - (-1)^{|X_i||X_j|}X_jX_i$.

Conjecture 10.7 *If $B = k\langle X_1, \dots, X_n\rangle/J$, where J is generated by r generic Hopf elements, then $B^!$ is a non-commutative c.i. or η is nilpotent, where $B = \mathrm{U}(\eta)$.*

Conjecture 10.8 *If $B = k\langle X_1, \dots, X_n\rangle/J$, where the variables are even and where J is generated by r generic Hopf elements, then $\eta(z) = (\log(1/(1-nz+rz^2)))_+$, where $B = \mathrm{U}(\eta)$.*

Finally we discuss the counterexamples to Conjecture 10.5. First we consider $(n, r) = (5, 7)$; i.e., 3 generic quadratic forms in 5 variables. It is shown in [Ei-Ko 94], that the forms $x_1x_4+x_2x_3, x_1x_5+x_2x_4, x_2x_5+x_3x_4$ are generic. The Lie algebra η for this case can be determined in the same way as in [Lö-Ro 97]. The result is that $\eta(z)$ is periodic, namely $\eta(z) = (5 + 3z)\prod_{i=0}^{\infty} z^{2i+1}$, while the expected series is $5z + 3z^2 + 5z^3$. In fact, η is the periodization of $sl(3)$ equipped with a $\mathbb{Z}_2$-grading. Secondly, if $(n, r) = (5, 8)$; i.e., 2 generic forms in 5 variables, the expected series is $5z + 2z^2$. Corollary 4.3 iii) together with Theorem 10.1 shows that $L^3 = \eta^3$ is 1-dimensional, so we really have a counterexample. We also have $L^4 = 0$, so in this case the Lie algebra is nilpotent but its series is not the expected minimal one.

11 Tables

We give the results of our calculations of $\eta_A(z)$ for the coordinate ring of s random points in P^n. For the definition of m and t, see Definition 5.3. For the definition of l_3, l_4, l_5, see Definition 6.3.

n	s	m	t	$\eta_A(z)$	l_3	l_4	l_5
3	7	1	3	$4z+3z^2$	7	7	7
3	6	2	2	$4z+4z^2+2z^3+3z^4+6z^5+\cdots$			
4	10	1	5	$5z+5z^2$	10	10	10
5	14	1	7	$6z+7z^2$	14	14	13
5	13	2	8	$6z+8z^2+5z^3+3z^4$			
6	19	1	9	$7z+9z^2$	18	18	17
6	18	2	4	$7z+10z^2+4z^3$			
6	17	2	10	$7z+11z^2+10z^3+10z^4$			
7	24	1	12	$8z+12z^2$	24	23	22
7	23	2	7	$8z+13z^2+7z^3$			
7	22	2	14	$8z+14z^2+14z^3+7z^4$			
7	21	2	21	$8z+15z^2+21z^3+42z^4+84z^5+91z^6$			
8	30	1	15	$9z+15z^2$	30	28	27
8	29	2	8	$9z+16z^2+8z^3$			
8	28	2	16	$9z+17z^2+16z^3$			
8	27	2	24	$9z+18z^2+24z^3+27z^4$			
8	26	2	32	$9z+19z^2+32z^3+73z^4+160z^5+176z^6$			
9	37	1	18	$10z+18z^2$	37	34	33
9	36	2	6	$10z+19z^2+6z^3$			
9	34	2	24	$10z+21z^2+24z^3$			
9	33	2	33	$10z+22z^2+33z^3+33z^4$			
9	32	2	42	$10z+23z^2+42z^3+91z^4+126z^5$			
10	44	1	22	$11z+22z^2$	44	41	39
10	43	2	10	$11z+23z^2+10z^3$			
10	41	2	30	$11z+25z^2+30z^3$			
10	40	2	40	$11z+26z^2+40z^3+10z^4$			
10	39	2	50	$11z+27z^2+50z^3+81z^4$			
10	38	2	60	$11z+28z^2+60z^3+153z^4+288z^5+\cdots$			
11	52	1	26	$12z+26z^2$	52	48	46
11	48	2	44	$12z+30z^2+44z^3$			
11	47	2	55	$12z+31z^2+55z^3+25z^4$			
11	46	2	66	$12z+32z^2+66z^3+111z^4$			
11	45	2	77	$12z+33z^2+77z^3+198z^4+\cdots$			

n	s	m	t	$\eta_A(z)$	l_3	l_4	l_5
12	61	1	30	$13z + 30z^2$	61	56	53
12	60	2	8	$13z + 31z^2 + 8z^3$			
12	56	2	56	$13z + 35z^2 + 56z^3$			
12	55	2	68	$13z + 36z^2 + 68z^3 + 3z^4$			
12	54	2	80	$13z + 37z^3 + 80z^3 + 105z^4$			
12	53	2	92	$13z + 38z^2 + 92z^3 + 208z^4 + \cdots$			
13	70	1	35	$14z + 35z^2$	70	64	61
13	64	2	78	$14z + 41z^2 + 78z^3$			
13	63	2	91	$14z + 42z^2 + 91z^3 + 42z^4$			
13	62	2	104	$14z + 43z^2 + 104z^3 + 162z^4$			
13	61	2	117	$14z + 44z^2 + 117z^3 + 283z^4 + \cdots$			
14	80	1	40	$15z + 40z^2$	80	73	70
14	73	2	98	$15z + 47z^2 + 98z^3$			
14	72	2	112	$15z + 48z^2 + 112z^3 + 36z^4$			
14	71	2	126	$15z + 49z^2 + 126z^3 + 175z^4 + \cdots$			
15	91	1	45	$16z + 45z^2$	91	82	79
15	90	2	10	$16z + 46z^2 + 10z^3$			
15	82	2	130	$16z + 54z^2 + 130z^3$			
16	102	1	51	$17z + 51z^2$	102	92	88
16	92	2	160	$17z + 61z^2 + 160z^3$			
17	114	1	57	$18z + 57z^2$	114	103	98
17	103	2	187	$18z + 68z^2 + 187z^3$			
18	127	1	63	$19z + 63z^2$	127	114	109
18	126	2	12	$19z + 64z^2 + 12z^3 + \cdots$			
19	140	1	70	$20z + 70z^2$	140	126	120
20	154	1	77	$21z + 77z^2$	154	139	132

Here are the results of our calculations of $\eta(z)$ for r random quadratic relations in an ordinary Lie algebra on n generators.

n	r	$\eta(z)$
4	5	$4z + z^2$
5	7	$5z + 3z^2 + 5z^3 + 3z^4 + 5z^5 + 3z^6 \cdots$
5	8	$5z + 2z^2 + z^3$
5	9	$5z + z^2$
6	10	$6z + 5z^2 + 10z^3 + 10z^4$
6	11	$6z + 4z^2 + 4z^3$
6	12	$6z + 3z^2$
7	13	$7z + 8z^2 + 21z^3 + 42z^4 + 84z^5 + 70z^6$
7	14	$7z + 7z^2 + 14z^3 + 7z^4$
7	15	$7z + 6z^2 + 7z^3$
7	16	$7z + 5z^2$
8	17	$8z + 11z^2 + 32z^3 + 73z^4 + 160z^5 + 144z^6$
8	18	$8z + 10z^2 + 24z^3 + 27z^4$
8	19	$8z + 9z^2 + 16z^3$
8	20	$8z + 8z^2 + 8z^3$
8	21	$8z + 7z^2$
9	22	$9z + 14z^2 + 42z^3 + 91z^4 + 126z^5$
9	23	$9z + 13z^2 + 33z^3$
9	24	$9z + 12z^2 + 24z^3$
9	25	$9z + 11z^2 + 15z^3$
9	26	$9z + 10z^2 + 6z^3$
9	27	$9z + 9z^2$

Referenser

[Ad 60] J.F. Adams, *On the non-existence of elements of Hopf invariant one*, Annals of Math. **72** (1960).

[An 82] D. Anick, *Noncommutative graded algebras and their Hilbert series*, J. Algebra **78** (1982), 120–140.

[Av 74] L. Avramov, *On the Hopf algebra of a local ring*, Isv. Akad. Nauk. SSSR, Ser. mat., **38** (1974), 253–277; English translation: Math. USSR, Izv. **8** (1974).

[Av 84] L. Avramov, *Local algebra and rational homotopy*, in: Homotopie Algébrique et Algèbre Locale, Astérisque 113/114 (1984) 15–43 .

[Ba-Ge 86] E. Ballico, A. V. Geramita, *The minimal free resolution of the ideal of s general points in P^3* , Proc. 1984 Vancouver conf. in alg. geom., 1–10, CMS Conf. Proc. **6**, AMS, Providence, 1986.

[Be-Kr 94] S. Beck, M. Kreuzer, *How to compute the canonical module of a set of points*, Algorithms in alg. geom. and appl. (Santander, 1994), Progr. Math **143**, 51–78, Birkhäuser, Basel, 1996.

[Bo 94] M. Boij, *Artin level algebras*, Dissertation KTH, Stockholm 1994.

[Ca-Ro-Va 91] M. P. Cavaliere, M. E. Rossi, G. Valla, *The Green-Lazarsfeld conjecture for $n+4$ points in P^n*, Rend. Sem. Mat. Univ. Pol. Torino **49** (1991), 175–195.

[Ch-Ei 48] C. Chevalley and S. Eilenberg, *Cohomology theory of Lie groups and Lie algebras*, Trans. Amer. Math. Soc. **63** (1948), 85–124.

[Co-Tr-Va 97] A. Conca, N. V. Trung, G. Valla, *Koszul property for points in projective space*, preprint 1997, acc. for publ. in Math. Scand.

[Ei-Ko 94] D. Eisenbud, J. Koh, *Nets of alternating matrices and the linear syzygy conjecture*, Adv. Math. **106** (1994), 1–35.

[Ei-Po 99] D. Eisenbud, S. Popescu, *Gale duality and free resolutions of ideals of points*, Inventiones Math. **136** (1999), 419–449.

[Fr 75] R. Fröberg, *Determination of a class of Poincaré series*, Math. Scand. **37** (1975), 29–39.

[Fr-Lö 91] R. Fröberg, C. Löfwall, *On Hilbert series for commutative and noncommutative graded algebras*, J. Pure Appl. Algebra **76** (1991), 33–38.

[Ge-Gr-Ro 86] A. Geramita, D. Gregory, L. Roberts, *Monomial ideals and points in projective space*, J. Pure Appl. Algebra **40** (1986), 33–62.

[Ge-Lo 89] A. V. Geramita, A. Lorenzini, *The Cohen-Macaulay type of $n+3$ points in P^n*, The Curves seminar at Queen's, vol. VI, no. F, Queen's Papers in Pure and Appl. Math. **83** (1989).

[Ge-Ma 84] A. V. Geramita, P. Maroscia, *The ideal of forms vanishing at a finite set of points in P^n*, J. Algebra **90** (1984), 528–555.

[Ge-Or 81] A. V. Geramita, F. Orecchia, *On the Cohen-Macaulay type of s lines in A^{n+1}*, J. Algebra **70** (1981), 116–140.

[Gu-La-St 91] V.K.A.M. Gugenheim, L. Lambe, J. Stasheff, Perturbation theory in differential homological algebra II, Il. J. Math., **35**(1991), 359–373.

[Gu 71] T. H. Gulliksen, *A homological characterization of complete intersections*, Comp. Math. **23** (1971), 251–255.

[Ha 92] S. Halperin, *Universal enveloping algebras and loop space homology*, J. Pure Appl. Algebra **83** (1992), 237–282.

[Hi-Si 96] A. Hirschowitz, C. Simpson, *La résolution minimale de l'idéal d'un arrangement général d'un grand nombre de points dans P^n* , Invent. Math. **126** (1996), 467–503.

[Ho-La 87] M. Hochster, D. Laksov, *The linear syzygies of generic forms*, Comm. Algebra **15** (1987), 227–239.

[Hu-Ka 91] J. Huebschmann, T. Kadeishvili, *Small models for chain algebras*, Math. Z. **207**, No.2 (1991), 245–280.

[Jo-La 00] L. Johansson, L. Lambe, *Transferring Algebra Structures up to Homology Equivalence*, to appear.

[La 97] F. Lauze, *Rang maximal pour T_{P^n}* , Manuscripta Math. **92** (1997), 525–543.

[Lo 93] A. Lorenzini, *The minimal resolution conjecture*, J. Algebra **156** (1993), 5–35.

[Lö 86] C. Löfwall, *On the subalgebra generated by the one-dimensional elements in the Yoneda Ext-algebra*, in: Algebra, Algebraic Topology and their Interactions, Lecture Notes in Math. **1183**, (Springer, Berlin 1986), 291–338.

[Lö 94] C. Löfwall, *Central elements and deformations of local rings*, J. Pure Appl. Algebra **91** (1994), 183–194.

[Lö-Ro 97] C. Löfwall, J.-E. Roos, *A nonnilpotent 1 - 2 -presented graded Hopf algebra whose Hilbert series converges in the unit circle*, Adv. Math. **130** (1997), 161–200.

[Ma 88] Y.I. Manin, *Quantum Groups and Non-Commutative Geometry*, Publ. CRM, Univ. de Montreal, 1988.

[Pr 70] X. Priddy, *Koszul resolution*, Trans. A.M.S. **152** (1970), 39–60.

[Wa 95] C. Walter, *The minimal free resolution of the homogeneous ideal of s general points*, Math. Z. **219** (1995), 231–234.