ISSN: 1401-5617

The ring of arithmetical functions with unitary convolution: Divisorial and topological properties

Jan Snellman

Research Reports in Mathematics
Number 4, 2002

Department of Mathematics
Stockholm University

Electronic versions of this document are available at
http://www.matematik.su.se/reports/2002/4

Date of publication: April 19, 2002
2000 Mathematics Subject Classification: Primary 11A25, Secondary 13J05.
Keywords: Unitary convolution, Schauder Basis, factorization.

Postal address:
Department of Mathematics
Stockholm University
S-106 91 Stockholm
Sweden

Electronic addresses:
http://www.matematik.su.se
info@matematik.su.se

THE RING OF ARITHMETICAL FUNCTIONS WITH UNITARY CONVOLUTION: DIVISORIAL AND TOPOLOGICAL PROPERTIES.

JAN SNELLMAN

Abstract. We study $(\mathcal{A}, +, \oplus)$, the ring of arithmetical functions with unitary convolution, giving an isomorphism between $(\mathcal{A}, +, \oplus)$ and a generalized power series ring on infinitely many variables, similar to the isomorphism of Cashwell-Everett[4] between the ring $(\mathcal{A}, +, \cdot)$ of arithmetical functions with *Dirichlet convolution* and the power series ring $\mathbb{C}[[x_1, x_2, x_3, \dots]]$ on countably many variables. We topologize it with respect to a natural norm, and shove that all ideals are quasi-finite. Some elementary results on factorization into atoms are obtained. We prove the existence of an abundance of non-associate regular non-units.

1. Introduction

The *ring of arithmetical functions with Dirichlet convolution*, which we'll denote by $(\mathcal{A}, +, \cdot)$, is the set of all functions $\mathbb{N}^+ \to \mathbb{C}$, where $\mathbb{N}^+$ denotes the positive integers. It is given the structure of a commutative $\mathbb{C}$-algebra by component-wise addition and multiplication by scalars, and by the Dirichlet convolution

$$f \cdot g(k) = \sum_{r|k} f(r)g(k/r). \tag{1}$$

Then, the multiplicative unit is the function e_1 with $e_1(1) = 1$ and $e_1(k) = 0$ for $k > 1$, and the additive unit is the zero function $\mathbf{0}$.

Cashwell-Everett [4] showed that $(\mathcal{A}, +, \cdot)$ is a UFD using the isomorphism

$$(\mathcal{A}, +, \cdot) \simeq \mathbb{C}[[x_1, x_2, x_3, \dots]], \tag{2}$$

where each x_i corresponds to the function which is 1 on the i'th prime number, and 0 otherwise.

Schwab and Silberberg [9] topologised $(\mathcal{A}, +, \cdot)$ by means of the norm

$$|f| = \frac{1}{\min\{\, k | f(k) \neq 0 \,\}} \tag{3}$$

They noted that this norm is an ultra-metric, and that $((\mathcal{A}, +, \cdot), |\cdot|)$ is a valued ring, i.e. that

(1) $|\mathbf{0}| = 0$ and $|f| > 0$ for $f \neq \mathbf{0}$,
(2) $|f - g| \leq \max\{|f|, |g|\}$,
(3) $|fg| = |f||g|$.

Date: April 19, 2002.
1991 *Mathematics Subject Classification.* 11A25, 13J05.
Key words and phrases. Unitary convolution, Schauder Basis, factorization.

They showed that $(\mathcal{A}, |\cdot|)$ is complete, and that each ideal is *quasi-finite*, which means that there exists a sequence $(e_k)_{k=1}^{\infty}$, with $|e_k| \to 0$, such that every element in the ideal can be written as a convergent sum $\sum_{k=1} c_k e_k$, with $c_k \in \mathcal{A}$.

In this article, we treat instead $(\mathcal{A}, +, \oplus)$, the ring of all arithmetical functions with unitary convolution. This ring has been studied by several authors, such as Vaidyanathaswamy [11], Cohen [5], and Yocom [13].

We topologise $\mathcal{A}$ in the same way as Schwab and Silberberg [9], so that $(\mathcal{A}, +, \oplus)$ becomes a normed ring (but, in contrast to $(\mathcal{A}, +, \cdot)$, not a valued ring). We show that all ideals in $(\mathcal{A}, +, \oplus)$ are quasi-finite.

We show that $(\mathcal{A}, +, \oplus)$ is isomorphic to a monomial quotient of a power series ring on countably many variables. It is présimplifiable and atomic, and there is a bound on the lengths of factorizations of a given element. We give a sufficient condition for nilpotency, and prove the existence of plenty of regular non-units.

Finally, we show that the set of arithmetical functions supported on square-free integers is a retract of $(\mathcal{A}, +, \oplus)$.

2. The ring of arithmetical functions with unitary convolution

Let p_i denote the i'th prime number, and denote by $\mathcal{P}$ the set of prime numbers. Let $\mathcal{PP}$ denote the set of prime powers. Let $\omega(r)$ denote the number of distinct prime factors of r, with $\omega(1) = 0$.

Definition 2.1. If k, m are positive integers, we define their *unitary product* as

$$k \oplus m = \begin{cases} km & \gcd(k, m) = 1 \\ 0 & \text{otherwise} \end{cases} \tag{4}$$

If $k \oplus m = p$, then we write $k \,||\, p$ and say that k is a *unitary divisor* of p.

The so-called *unitary convolution* was introduced by Vaidyanathaswamy [11], and was further studied Eckford Cohen [5].

Definition 2.2. $\mathcal{A} = \{f : \mathbb{N}^+ \to \mathbb{C}\}$, the set of complex-valued functions on the positive integers. We define the *unitary convolution* of $f, g \in \mathcal{A}$ as

$$(f \oplus g)(n) = \sum_{\substack{m \oplus p = n \\ m,n \geq 1}} f(m)g(n) = \sum_{d \,||\, n} f(d)g(n/d) \tag{5}$$

and the addition as

$$(f + g)(n) = f(n) + g(n)$$

The ring $(\mathcal{A}, +, \oplus)$ is called *the ring of arithmetic functions* with unitary convolution.

Definition 2.3. For each positive integer k, we define $e_k \in \mathcal{A}$ by

$$e_k(n) = \begin{cases} 1 & k = n \\ 0 & k \neq n \end{cases} \tag{6}$$

We also define[1] $\mathbf{0}$ as the zero function, and $\mathbf{1}$ as the function which is constantly 1.

Lemma 2.4. *$\mathbf{0}$ is the additive unit of $\mathcal{A}$, and e_1 is the multiplicative unit. We have that*

$$(e_{k_1} \oplus e_{k_2} \oplus \cdots \oplus e_{k_r})(n) = \begin{cases} 1 & n = k_1 k_2 \cdots k_r \text{ and } \gcd(k_i, k_j) = 1 \text{ for } i \neq j \\ 0 & \text{otherwise} \end{cases} \tag{7}$$

hence

$$e_{k_1} \oplus e_{k_2} \oplus \cdots \oplus e_{k_r} = \begin{cases} e_{k_1 k_2 \cdots k_r} & \text{if } \gcd(k_i, k_j) = 1 \text{ for } i \neq j \\ 0 & \text{otherwise} \end{cases} \tag{8}$$

Proof. The first assertions are trivial. We have [10] that for $f_1, \ldots, f_r \in \mathcal{A}$,

$$(f_1 \oplus \cdots f_r)(n) = \sum_{a_1 \oplus \cdots a_r = n} f_1(a_1) \cdots f_r(a_r) \tag{9}$$

Since

$$e_{k_1}(a_1) e_{k_2}(a_2) \cdots e_{k_r})(a_r) = 1 \text{ iff } \forall i : k_i = a_i,$$

(7) follows. □

Lemma 2.5. *Any e_n can be uniquely expressed as a square-free monomial in $\{ e_k | k \in \mathcal{PP} \}$.*

Proof. By unique factorization, there is a unique way of writing $n = p_{i_1}^{a_1} \cdots p_{i_r}^{a_r}$, and (8) gives that

$$e_n = e_{p_{i_1}^{a_1} \cdots p_{i_r}^{a_r}} = e_{p_{i_1}^{a_1}} \oplus \cdots e_{p_{i_r}^{a_r}}.$$

□

Theorem 2.6. *$(\mathcal{A}, +, \oplus)$ is a quasi-local, non-noetherian commutative ring having divisors of zero. The units $U(\mathcal{A})$ consists of those f such that $f(1) \neq 0$.*

Proof. It is shown in [10] that $(\mathcal{A}, +, \oplus)$ is a commutative ring, having zero-divisors, and that the units consists of those f such that $f(1) \neq 0$. If $f(1) = 0$ then

$$(f \oplus g)(1) = f(1)g(1) = 0.$$

Hence the non-units form an ideal $\mathfrak{m}$, which is then the unique maximal ideal.

We will show (Lemma 3.10) that $\mathfrak{m}$ contains an ideal (the ideal generated by all e_k, for $k > 1$) which is not finitely generated, so $\mathcal{A}$ is non-noetherian. □

[1] In [10], $\mathbf{1}$ is denoted e, and e_1 denoted e_0.

3. A topology on $\mathcal{A}$

The results of this section are inspired by [9], were the authors studied the ring of arithmetical functions under Dirichlet convolution. We'll use the notations of [3]. We regard $\mathbb{C}$ as trivially normed.

Definition 3.1. Let $f \in \mathcal{A} \setminus \{\mathbf{0}\}$. We define the *support* of f as

$$\operatorname{supp}(f) = \left\{ n \in \mathbb{N}^+ \middle| f(n) \neq 0 \right\} \tag{10}$$

We define the *order*[2] of a non-zero element by

$$\mathrm{N}(f) = \min \operatorname{supp}(f) \tag{11}$$

We also define the *norm* of f as

$$|f| = \mathrm{N}(f)^{-1} \tag{12}$$

and the *degree* as

$$\mathrm{D}(f) = \min \left\{ \omega(k) \middle| k \in \operatorname{supp}(f) \right\} \tag{13}$$

By definition, the zero element has order infinity, norm 0, and degree -1.

Lemma 3.2. *The value semigroup of* $(\mathcal{A}, |\cdot|)$ *is* $|\mathcal{A} \setminus \{\mathbf{0}\}| = \left\{ 1/k \middle| k \in \mathbb{N}^+ \right\}$, *a discrete subset of* $\mathbb{R}^+$.

Lemma 3.3. *Let* $f, g \in \mathcal{A} \setminus \{\mathbf{0}\}$. *Let* $\mathrm{N}(f) = i$, $\mathrm{N}(g) = j$, *so that* $f(i) \neq 0$ *but* $f(k) = 0$ *for all* $k < i$, *and similarly for* g. *We assume that* $i \leq j$. *Then, the following hold:*

(i) $\mathrm{N}(f - g) \geq \min \{\mathrm{N}(f), \mathrm{N}(g)\}$.
(ii) $\mathrm{N}(cf) = \mathrm{N}(f)$ *for* $c \in \mathbb{C} \setminus \{0\}$.
(iii) $\mathrm{N}(f) = 1$ *iff* f *is a unit.*
(iv) $\mathrm{N}(f \cdot g) = \mathrm{N}(f)\mathrm{N}(g) \leq \mathrm{N}(f \oplus g)$, *with equality iff* $(i, j) = 1$.
(v) $\mathrm{N}(f \oplus g) \geq \max \{\mathrm{N}(f), \mathrm{N}(g)\}$, *with strict inequality iff both* f *and* g *are non-units.*
(vi) $\mathrm{D}(f + g) \geq \min \mathrm{D}(f), \mathrm{D}(g)$.
(vii) $\mathrm{D}(f \cdot g) = \mathrm{D}(f) + \mathrm{D}(g)$.
(viii) $\mathrm{D}(f) = 0$ *if and only if* f *is a unit.*
(ix) *Suppose that* $f \oplus g \neq \mathbf{0}$. *Then*

$$\mathrm{D}(f \oplus g) \geq \mathrm{D}(f) + \mathrm{D}(g) \geq \max \{\mathrm{D}(f), \mathrm{D}(g)\}.$$

with $\mathrm{D}(f) + \mathrm{D}(g) > \max \{\mathrm{D}(f), \mathrm{D}(g)\}$ *if* f, g *are non-units.*

Proof. (i), (ii), and (iii) are trivial, and (iv) is proved in [10]. (vi), (vii), and (viii) are proved in [8]. Let m be a monomial in the support of f such that $\mathrm{D}(m) = \mathrm{D}(f)$, and let n be a monomial in the support of g such that $\mathrm{D}(n) = \mathrm{D}(g)$. For any a in the support of f and any q in the support of g, such that $a \oplus q \neq 0$, we have that

$$\mathrm{D}(a \oplus q) = \mathrm{D}(a) + \mathrm{D}(q) \geq \mathrm{D}(f) + \mathrm{D}(g).$$

This proves (ix). (v) is proved similarly. □

Corollary 3.4. $|f \oplus g| \leq |f||g| = |f \cdot g|$.

[2] In [10] the term *norm* is used.

Proposition 3.5. *$|\cdot|$ is an ultrametric function on $\mathcal{A}$, making $(\mathcal{A}, +, \oplus)$ a normed ring, as well as a faithfully normed, b-separable complete vector space over $\mathbb{C}$.*

Proof. $((\mathcal{A}, +, \cdot), |\cdot|)$ is a valuated ring, and a faithfully normed complete vector space over $\mathbb{C}$ [9]. It is also separable with respect to bounded maps [3, Corollary 2.2.3]. So $(\mathcal{A}, +)$ is a normed group, hence Corollary 3.4 shows that $(\mathcal{A}, +, \oplus)$ is a normed ring. □

Note that, unlike $((\mathcal{A}, +, \cdot), |\cdot|)$, the normed ring $((\mathcal{A}, +, \oplus), |\cdot|)$ is not a valued ring, since

$$|e_2 \oplus e_2| = |\mathbf{0}| = 0 < |e_2|^2 = 1/4.$$

In fact, we have that

Lemma 3.6. *If f is a unit, then $1 = |f^n| = |f|^n$ for all positive integers n. If n is a non-unit, then $|f^n| < |f|^n$ for all $n > 1$.*

Proof. The first assertion is trivial, so suppose that f is a non-unit. From Corollary 3.4 we have that $|f^n| \leq |f|^n$. If $|f| = 1/k$, $k > 1$, i.e. $f(k) \neq 0$ but $f(j) = 0$ for $j < k$, then $f^2(k^2) = 0$ since $\gcd(k, k) = k > 1$. It follows that $\left|f^2\right| > |f|^2$, from which the result follows. □

Recall that in a normed ring, a non-zero element f is called

- *topologically nilpotent* if $f^n \to 0$,
- *power-multiplicative* if $|f^n| = |f|^n$ for all n,
- *multiplicative* if $|fg| = |f||g|$ for all g in the ring.

Theorem 3.7. *Let $f \in ((\mathcal{A}, +, \oplus), |\cdot|)$, $f \neq \mathbf{0}$. Then the following are equivalent:*

(1) f is topologically nilpotent,
(2) f is not power-multiplicative,
(3) f is not multiplicative[3] in the normed ring $(\mathcal{A}, +, \oplus), |\cdot|)$,
(4) f is a non-unit,
(5) $|f| < 1$.

Proof. Using [3, 1.2.2, Prop. 2], this follows from the previous Lemma, and the fact that for a unit f,

$$1 = \left|f^{-1}\right| = |f|^{-1}.$$

□

3.1. A Schauder basis for $(\mathcal{A}, |\cdot|)$.

Definition 3.8. Let $\mathcal{A}'$ denote the subset of $\mathcal{A}$ consisting of functions with finite support. We define a pairing

$$\begin{aligned} \mathcal{A} \times \mathcal{A}' &\to \mathbb{C} \\ \langle f, g \rangle &= \sum_{k=1}^{\infty} f(k) g(k) \end{aligned} \tag{14}$$

[3]This is not the same concept as multiplicativity for arithmetical functions, i.e. that $f(nm) = f(n)f(m)$ whenever $(n, m) = 1$. However, since the latter kind of elements satisfy $f(1) = 1$, they are units, and hence multiplicative in the normed-ring sense.

Theorem 3.9. *The set* $\{ e_k | k \in \mathbb{N}^+ \}$ *is an ordered orthogonal Schauder base in the normed vector space* $(\mathcal{A}, |\cdot|)$. *In other words, if* $f \in \mathcal{A}$ *then*

$$f = \sum_{k=1}^{\infty} c_k e_k, \qquad c_k \in \mathbb{C} \tag{15}$$

where

(i) $|e_k| \to 0$,
(ii) the infinite sum (15) *converges w.r.t. the ultrametric topology,*
(iii) the coefficients c_k *are uniquely determined by the fact that*

$$\langle f, e_k \rangle = f(k) = c_k \tag{16}$$

(iv)

$$\max_{k \in \mathbb{N}^+} \{|c_k||e_k|\} = \left| \sum_{k=1}^{\infty} c_k e_k \right| \tag{17}$$

The set $\{ e_p | p \in \mathcal{PP} \}$ *generates a dense subalgebra of* $((\mathcal{A}, +, \oplus), |\cdot|)$.

Proof. It is proved in [9] that this set is a Schauder base in the topological vector space $(\mathcal{A}, |\cdot|)$. It also follows from [9] that the coefficients c_k in (3.9) are given by $c_k = f(k)$.

It remains to prove orthogonality. With the above notation,

$$|f| = \left| \sum_{k=1}^{\infty} c_k e_k \right| = 1/j,$$

where j is the smallest k such that $c_k \neq 0$. Recalling that $\mathbb{C}$ is trivially normed, we have that

$$|c_k||e_k| = \begin{cases} |e_k| = 1/k & \text{if } c_k \neq 0 \\ 0 & \text{if } c_k = 0 \end{cases},$$

so $\max_{k \in \mathbb{N}^+} \{|c_k||e_k|\} = 1/j$, with j as above, so (17) holds.

By Lemma 2.5 any e_k can be written as a square-free monomial in the elements of $\{ e_p | p \in \mathcal{PP} \}$. The set $\{ e_k | k \in \mathbb{N}^+ \}$ is dense in $\mathcal{A}$, so $\{ e_p | p \in \mathcal{PP} \}$ generates a dense subalgebra. □

Let $J \subset \mathfrak{m}$ denote the ideal generated by all e_k, $k > 1$.

Lemma 3.10. *J is not finitely generated.*

Proof. If J is finitely generated, then there is an N such that

$$J = (e_2, \ldots, e_N).$$

Let L be a prime number, $L > N$. Since $e_L \in J$, we have that

$$e_L = \sum_{k=2}^{N} f_k \oplus e_k, \qquad f_k \in \mathcal{A}.$$

We write $f_k = \sum_{i=1}^{\infty} c_{ki} e_i$, so that

$$e_L = \sum_{k=2}^{N} e_k \oplus \sum_{i=1}^{\infty} c_{ik} e_i = \sum_{k=2}^{N} \sum_{i=1}^{\infty} c_{ik} e_i \oplus e_k = \sum_{k=2}^{N} \sum_{\gcd(i,k)=1} c_{ik} e_{ik}.$$

But this is impossible, because we can not write $L = ik$ with $\gcd(i,k) = 1$ and $2 \leq i \leq N < L$. □

Definition 3.11. An ideal $I \subset \mathcal{A}$ is called quasi-finite if there exists a sequence $(g_k)_{k=1}^{\infty}$ in I such that $|g_k| \to 0$ and such that every element $f \in I$ can be written (not necessarily uniquely) as a convergent sum

$$f = \sum_{k=0}^{\infty} a_k \oplus g_k, \qquad a_k \in \mathcal{A} \tag{18}$$

Lemma 3.12. $\mathfrak{m}$ *is quasi-finite.*

Proof. By Theorem 3.9 the set $\{\, e_k | k > 1 \,\}$ is a quasi-finite generating set for $\mathfrak{m}$. □

Since all ideals are contained in $\mathfrak{m}$, it follows that any ideal containing $\{\, e_k | k > 1 \,\}$ is quasi-finite. Furthermore, such an ideal has $\mathfrak{m}$ as its closure. In particular, J is quasi-finite, but not closed.

Theorem 3.13. *All (non-zero) ideals in $\mathcal{A}$ are quasi-finite. In fact, given any subspace I if we can find*

$$G(I) := (g_k)_{k=1}^{\infty} \tag{19}$$

such that for all $f \in I$,

$$\exists c_1, c_2, c_3, \cdots \in \mathbb{C}, \qquad f = \sum_{i=1}^{\infty} c_i g_i. \tag{20}$$

So all subspaces possesses a Schauder basis.

Proof. We construct $G(I)$ in the following way: for each

$$k \in \{\, \mathrm{N}(f) \,|\, f \in I \setminus \{\mathbf{0}\} \,\} =: N(I)$$

we choose a $g_k \in I$ with $\mathrm{N}(g_k) = k$, and with $g_k(k) = 1$. In other words, we make sure that the "leading coefficient" is 1; this can always be achieved since the coefficients lie in a field. For $k \notin N(I)$ we put $g_k = \mathbf{0}$.

To show that this choice of elements satisfy (20), take any $f \in I$, and put $f_0 = f$. Then define recursively, as long as $f_i \neq \mathbf{0}$,

$$\begin{aligned} n_i &:= N(f_i) \\ \mathbb{C} \ni a_i &:= f_i(n_i) \\ \mathcal{A} \ni f_{i+1} &:= f_i - a_i g_{n_i} \end{aligned}$$

Of course, if $f_i = \mathbf{0}$, then we have expressed f as a linear combination of

$$g_{n_1}, \ldots, g_{n_{i-1}},$$

and we are done. Otherwise, note that by induction $f_i \in I$, so $n_i \in N(I)$, hence $g_{n_i} \neq 0$. Thus $\mathrm{N}(f_{i+1}) > \mathrm{N}(f_i)$, so $|f_{i+1}| < |f_i|$, whence

$$|f_0| > |f_1| > |f_2| > \cdots \to 0.$$

But

$$f_{i+1} = f - \sum_{j=1}^{i} a_j g_{n_j},$$

so

$$F_i := \sum_{j=1}^{i} a_j g_{n_j} \to f,$$

which shows that $\sum_{j=1}^{\infty} a_j g_j = f$. □

4. A fundamental isomorphism

4.1. **The monoid of separated monomials.** Let

$$Y = \left\{ y_i^{(j)} \middle| i, j \in \mathbb{N}^+ \right\} \tag{21}$$

be an infinite set of variables, in bijective correspondence with the integer lattice points in the first quadrant minus the axes. We call the subset

$$Y_i = \left\{ y_i^{(j)} \middle| j \in \mathbb{N}^+ \right\} \tag{22}$$

the *i'th column* of Y.

Let $[Y]$ denote the free abelian monoid on Y, and let $\mathcal{M}$ be the subset of *separated monomials*, i.e. monomials in which no two occurring variables come from the same column:

$$\mathcal{M} = \left\{ y_{i_1}^{(j_1)} y_{i_2}^{(j_1)} \cdots y_{i_r}^{(j_r)} \middle| 1 \leq i_i < i_2 < \cdots i_r \right\} \tag{23}$$

We regard $\mathcal{M}$ as a monoid-with-zero, so that the multiplication is given by

$$m \oplus m' = \begin{cases} mm' & mm' \in \mathcal{M} \\ 0 & \text{otherwise} \end{cases} \tag{24}$$

Note that the zero is exterior to $\mathcal{M}$, i.e. $0 \notin \mathcal{M}$. The set $\mathcal{M} \cup \{0\}$ is a (non-cancellative) monoid if we define $m \oplus 0 = 0$ for all $m \in \mathcal{M}$.

Recall that $\mathcal{PP}$ denotes the set of prime powers. It follows from the fundamental theorem of arithmetic that any positive integer n can be uniquely written as a *square-free* product of prime powers. Hence we have that

$$\begin{aligned} \Phi : Y &\to \mathcal{PP} \\ y_i^{(j)} &\mapsto p_i^j \end{aligned} \tag{25}$$

is a bijection which can be extended to a bijection

$$\begin{aligned} \Phi : \mathcal{M} &\to \mathbb{N}^+ \\ 1 &\mapsto 1 \\ y_{i_1}^{(j_1)} \cdots y_{i_r}^{(j_r)} &\mapsto p_{i_1}^{j_1} \cdots p_{i_r}^{j_r} \end{aligned} \tag{26}$$

If we regard $\mathbb{N}^+$ as a monoid-with-zero with the operation $\oplus$ of (4), then (26) is a monoid-with-zero isomorphism.

4.2. **The ring $\mathcal{A}$ as a generalized power series ring, and as a quotient of $\mathbb{C}[[Y]]$.** Let R be the large power series ring on $[Y]$, i.e. $R = C[[Y]]$ consists of all formal power series $\sum c_{\boldsymbol{\alpha}} \boldsymbol{y}^{\boldsymbol{\alpha}}$, where the sum is over all multisets $\boldsymbol{\alpha}$ on Y.

Let S be the generalized monoid-with-zero ring on $\mathcal{M}$. By this, we mean that S is the set of all formal power series

$$\sum_{m \in \mathcal{M}} f(m)m \tag{27}$$

with component-wise addition, and with multiplication

$$\begin{gathered}\left(\sum_{m \in \mathcal{M}} f(m)m\right) \oplus \left(\sum_{m \in \mathcal{M}} g(m)m\right) = \left(\sum_{m \in \mathcal{M}} h(m)m\right) \\ h(m) = (f \oplus g)(m) = \sum_{s \oplus t = m} f(s)g(t)\end{gathered} \tag{28}$$

Define

$$\operatorname{supp}(\sum_{m \in [Y]} c_m m) = \{\, m \in Y \,|\, c_m \neq 0 \,\} \tag{29}$$

$$\operatorname{supp}(\sum_{m \in \mathcal{M}} c_m m) = \{\, m \in \mathcal{M} \,|\, c_m \neq 0 \,\} \tag{30}$$

$$\tag{31}$$

Let furthermore

$$\mathfrak{D} = \{\, f \in R \,|\, \operatorname{supp}(f) \cap \mathcal{M} = \emptyset \,\} \tag{32}$$

Theorem 4.1. *S and $\frac{R}{\mathfrak{D}}$ and $\mathcal{A}$ are isomorphic as $\mathbb{C}$-algebras.*

Proof. The bijection (26) induces a bijection between S and $\mathcal{A}$ which is an isomorphism because of the way multiplication is defined on S. In detail, the isomorphism is defined by

$$\begin{gathered} S \ni \sum_{m \in \mathcal{M}} c_m m \mapsto f \in \mathcal{A} \\ f(\Phi(m)) = c_m \end{gathered} \tag{33}$$

For the second part, consider the epimorphism

$$\phi : R \to S$$

$$\phi\left(\sum_{m \in [Y]} c_m m\right) = \sum_{m \in \mathcal{M}} c_m m$$

Clearly, $\ker(\phi) = \mathfrak{D}$, hence $S \simeq \frac{R}{\ker(\phi)} = \frac{R}{\mathfrak{D}}$. □

Let us exemplify this isomorphism by noting that e_n, where n has the square-free factorization $n = p_1^{a_1} \cdots p_r^{a_r}$, corresponds to the square-free monomial $y_1^{(a_1)} \cdots y_r^{(a_r)}$, and that

$$\mathbf{1} = \sum_{m \in \mathcal{M}} m = \prod_{i=1}^{\infty} \left(1 + \sum_{j=1}^{\infty} y_i^{(j)}\right) \tag{34}$$

What does its inverse μ^* correspond to?

Definition 4.2. For $m \in \mathcal{M}$, we denote by $\mathrm{D}(m)$ the number of occurring variables in m (by definition, $\mathrm{D}(1) = 0$ and $\mathrm{D}(0) = -\infty$). For

$$S \ni f = \sum_{m \in \mathcal{M}} c_m m$$

we put

$$\mathrm{D}(f) = \min \{ \mathrm{D}(m) | c_m \neq 0 \} \tag{35}$$

Using the isomorphism between S and $\mathcal{A}$, we define $\mathrm{D}(g)$ for any $g \in \mathcal{A}$ by

$$\mathrm{D}(g) = \min \{ \omega(n) | f(n) \neq 0 \} .$$

It is known (see [10]) that

$$\mu^*(r) = (-1)^{\omega(r)} \tag{36}$$

We then have that μ^* corresponds to

$$\mathbf{1}^{-1} = \frac{1}{\prod_{i=1}^{\infty} \left(1 + \sum_{j=1}^{\infty} y_i^{(j)}\right)} = \prod_{i=1}^{\infty} \frac{1}{1 + \sum_{j=1}^{\infty} y_i^{(j)}} = \sum_{m \in \mathcal{M}} (-1)^{\mathrm{D}(m)} m \tag{37}$$

Recall that $f \in \mathcal{A}$ is a *multiplicative* arithmetic function if $f(nm) = f(n)f(m)$ whenever $(n, m) = 1$. Regarding f as an element of S we have that f is multiplicative if and only if it can be written as

$$f = \prod_{i=1}^{\infty} \left(1 + \sum_{j=1}^{\infty} c_{i,j} y_i^{(j)} \right) \tag{38}$$

It is now easy to see that the multiplicative functions form a group under multiplication.

4.3. **The continuous endomorphisms.** In [9], Schwab and Silberberg characterized all continuous endomorphisms of Γ. We give the corresponding result for $\mathcal{A}$:

Theorem 4.3. *Every continuous endomorphism θ of the $\mathbb{C}$-algebra $S \simeq \mathcal{A}$ is defined by*

$$\theta(y_i^{(j)}) = \gamma_{i,j} \tag{39}$$

where

$$\gamma_{i,j}\gamma_{i,k} = 0 \qquad \textit{for all } i, j, k \tag{40}$$

and

$$\gamma_{a_1(n),b_1(n)} \cdots \gamma_{a_r(n),b_r(n)} \to 0 \qquad \textit{as } n = p_{a_1(n)}^{b_1(n)} \cdots p_{a_r(n)}^{b_r(n)} \to \infty \tag{41}$$

Proof. Recall that $S \simeq \frac{R}{\mathfrak{D}}$, where $R = \mathbb{C}[[Y]]$ and $\mathfrak{D}$ is the closure of the ideal generated by all non-separated quadratic monomials $y_i^{(j)} y_i^{(k)}$. Since the set of square-free monomials in the $y_i^{(j)}$'s form a Schauder base, any continuous C-algebra endomorphism θ of S is determined by its values on the $y_i^{(j)}$'s, and must fulfill (41). Since $y_i^{(j)} y_i^{(k)} = 0$ in S, we must have that

$$\theta(0) = \theta(y_i^{(j)} y_i^{(k)}) = \theta(y_i^{(j)})\theta(y_i^{(k)}) = \gamma_{i,j}\gamma_{i,k} = 0.$$

□

5. Nilpotent elements and zero divisors

Definition 5.1. For $m \in \mathbb{N}^+$, define the *prime support* of m as

$$\operatorname{psupp}(m) = \{\, p \in \mathcal{P} \,|\, p \,|\, m \,\} \tag{42}$$

and (when $m > 1$) the *leading prime* as

$$\operatorname{lp}(m) = \min \operatorname{psupp}(m) \tag{43}$$

For $n \in \mathbb{N}^+$, put

$$A^n = \left\{\, k \in \mathbb{N}^+ \middle| p_n \,|\, k \text{ but } p_i \not| k \text{ for } i < n \,\right\} = \left\{\, k \in \mathbb{N}^+ \middle| \operatorname{lp}(k) = p_n \,\right\} \tag{44}$$

Then $\mathbb{N}^+ \setminus \{1\}$ is a disjoint union

$$\mathbb{N}^+ \setminus \{1\} = \bigsqcup_{i=1}^{\infty} A^i \tag{45}$$

Definition 5.2. Let $f \in \mathcal{A}$ be a non-unit. The *canonical decomposition* of f is the unique way of expressing f as a convergent sum

$$f = \sum_{i=1}^{\infty} f_i, \quad f_i = \sum_{k \in A^i} f(k) e_k \tag{46}$$

The element f is said to be of *polynomial type* if all but finitely many of the f_i's are zero. In that case, the largest N such that $f_N \neq \mathbf{0}$ is called the *filtration degree* of f.

Lemma 5.3.

$$f_i = \sum_{j=1}^{\infty} e_{p_i^j} \oplus g_{i,j}, \quad r \leq i, \quad p_r \,|\, n \implies g_{i,j}(n) = 0. \tag{47}$$

For any n there is at most one pair (i,j) such that

$$\left(e_{p_i^j} \oplus g_{i,j} \right)(n) \neq 0.$$

More precisely, if

$$n = p_{i_1}^{j_1} \cdots p_{i_r}^{j_r}, \qquad i_1 < \cdots < i_r,$$

then $\left(e_{p_{i_1}^{j_1}} \oplus g_{i_1,j_1} \right)(n)$ *may be non zero.*

Definition 5.4. For $k \in \mathbb{N}$, define

$$I_k = \left\{\, f \in \mathcal{A} \middle| f(n) = 0 \text{ for every } n \text{ such that } (n, p_1 p_2 \cdots p_k) = 1 \,\right\} \tag{48}$$

Lemma 5.5. *I_k is an ideal in $(\mathcal{A}, +, \oplus)$.*

Proof. It is shown in [8] that the I_k's form an ascending chain of ideals in $(\mathcal{A}, +, \cdot)$. They are also easily seen to be ideals in $(\mathcal{A}, +, \oplus)$: if

$$f \in I_k,\ g \in \mathcal{A} \text{ and } (n, p_1 p_2 \cdots p_k) = 1$$

then

$$(f \oplus g)(n) = \sum_{d \| n} f(d) g(n/d) = 0,$$

since $(d, p_1 p_2 \cdots p_k) = 1$ for any unitary divisor of n. $\square$

Theorem 5.6. *Let $N \in \mathbb{N}^+$, and let $f \in (\mathcal{A}, +, \oplus)$ be a non-unit. Then*

$$\begin{aligned} I_N &= \operatorname{ann}(e_{p_1 \cdots p_N}) \\ &= \{\mathbf{0}\} \cup \{\, f \in \mathcal{A} | f \text{ is of polynomial type and has filtration degree } N \,\} \\ &= \overline{\mathcal{A}\left\{\, e_{p_i^a} \middle| a, i \in \mathbb{N}^+,\, i \leq N \,\right\}} \end{aligned}$$

where $\overline{\mathcal{A}W}$ denotes the topological closure of the ideal generated by the set W.

Proof. If $f \in I_N$ then for all k

$$(f \oplus e_{p_1 \cdots p_N})(k) = \sum_{a \oplus p_1 \cdots p_N = k} f(a) e_{p_1 \cdots p_N}(p_1 \cdots p_N) = \sum_{a \oplus p_1 \cdots p_N = k} f(a) = 0 \tag{49}$$

so $f \in \operatorname{ann}(e_{p_1 \cdots p_N})$. Conversely, if $f \in \operatorname{ann}(e_{p_1 \cdots p_N})$ then $(f \oplus e_{p_1 \cdots p_N})(k) = 0$ for all k, hence if $(n, p_1 \cdots p_N) = 1$ then

$$0 = (f \oplus e_{p_1 \cdots p_N})(n p_1 \cdots p_N) = f(n) e_{p_1 \cdots p_N}(p_1 \cdots p_N) = f(n) \tag{50}$$

hence $f \in I_N$.

If $f \in I_N$ then for $j > N$ we get that $f_j = \mathbf{0}$, since

$$f_j(k) = \begin{cases} 0 & \text{if } k \notin A^j \\ f(k) = 0 & \text{if } k \in A^j \end{cases}$$

Hence $f = \sum_{i=1}^N f_i$. Conversely, if f can be expressed thusly, then $f(k) = f_{j_1}(k) = 0$ for $k = p_{j_1}^{a_1} \cdots p_{j_r}^{a_r}$ with $N < j_1 < \cdots < j_r$.

The last equality follows from Theorem 3.9. □

Theorem 5.7. *Let $f \in \mathcal{A}$ be a non-unit. The following are equivalent:*

(i) f is of polynomial type.
(ii) $f \in \cup_{k=0}^{\infty} I_k$,
(iii) There is a finite subset $Q \subset \mathcal{P}$ such that $f(k) = 0$ for all k relatively prime to all $p \in Q$.
(iv) $f \in \cup_{N=1}^{\infty} \operatorname{ann}(e_{p_1 p_2 \cdots p_N})$.
(v) f is contained in the topological closure of the ideal generated by the set $\left\{\, e_{p_i^a} \middle| a, i \in \mathbb{N}^+,\, i \leq N \,\right\}$.

If f has finite support, then it is of polynomial type. If f is of polynomial type, then it is nilpotent.

Proof. Clearly, a finitely supported f is of polynomial type. The equivalence (i) $\iff$ (ii) $\iff$ (iii) $\iff$ (iv) $\iff$ (v) follows from the previous theorem.

If f is of polynomial type, say of filtration degree N, then

$$f = \sum_{i=1}^N f_i \tag{51}$$

and we see that if f^{N+1} is the $N+1$'st unitary power of f, then f^{N+1} is the linear combination of monomials in the f_i's, and none of these monomials are square-free. Since $f_i \oplus f_i = \mathbf{0}$ for all i, we have that $f^{N+1} = \mathbf{0}$. So f is nilpotent. □

Lemma 5.8. *The elements of polynomial type forms an ideal.*

Proof. By the previous theorem, this set can be expressed as

$$\bigcup_{n=1}^{\infty} I_n,$$

which is an ideal since each I_n is. □

Question 5.9. *Are all [nilpotent elements, zero divisors] of polynomial type? If one could prove that the zero divisors are precisely the elements of polynomial type, then by Lemma 5.8 it would follow that $Z(\mathcal{A})$ is an ideal, and moreover a prime ideal, since the product of two regular elements is regular (in any commutative ring). Then one could conclude* [6] *that $(\mathcal{A}, +, \oplus)$ has* few zero divisors*, hence is* additively regular*, hence is a* Marot ring.

Theorem 5.10. *$(\mathcal{A}, +, \oplus)$ contains infinitely many non-associate regular non-units.*

Proof. **Step 1**. We first show that there is at least one such element. Let $f \in \mathcal{A}$ denote the arithmetical function

$$f(k) = \begin{cases} 1 & k \in \mathcal{PP} \\ 0 & \text{otherwise} \end{cases}$$

Then f is a non-unit, and using a result by Yocom [13, 8] we have that f is contained in a subring of $(\mathcal{A}, +, \oplus)$ which is a discrete valuation ring isomorphic to $\mathbb{C}[[t]]$, the power series ring in one indeterminate. This ring is a domain, so f is not nilpotent.

We claim that f is in fact regular. To show this, suppose that $g \in \mathcal{A}$, $f \oplus g = \mathbf{0}$. We will show that $g = \mathbf{0}$.

Any positive integer m can be written $m = q_1^{a_1} \cdots q_r^{a_r}$, where the q_i are distinct prime numbers. If $r = 0$, then $m = 1$, and $g(1) = 0$, since

$$0 = (f \oplus g)(2) = f(2)g(1) = g(1).$$

For the case $r = 1$, we want to show that $g(q^a) = 0$ for all prime numbers q. Choose three different prime powers $q_1^{a_1}$, $q_2^{a_2}$, and $q_3^{a_3}$. Then

$$0 = f \oplus g(q_i^{a_i} q_j^{a_j}) = f(q_i^{a_i})g(q_j^{a_j}) + f(q_j^{a_j})g(q_i^{a_i}) = g(q_j^{a_j}) + g(q_i^{a_i}),$$

when $i \neq j$, $i, j \in \{1, 2, 3\}$. In matrix notation, these three equations can be written as

$$\begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} g(q_1^{a_1}) \\ g(q_2^{a_2}) \\ g(q_3^{a_3}) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

from which we conclude (since the determinant of the coefficient matrix is non-zero) that $0 = g(q_1^{a_1}) = g(q_2^{a_2}) = g(q_3^{a_3})$.

Now for the general case, $r > 1$. We need to show that that

$$g(q_1^{a_1} \cdots q_r^{a_r}) = 0 \tag{52}$$

whenever $q_1^{a_1}, \ldots, q_r^{a_r}$ are pair-wise relatively prime prime powers.

Choose N pair-wise relatively prime prime powers $q_1^{a_1}, \ldots, q_N^{a_N}$. For each $r+1$-subset $q_{s_1}, \ldots, q_{s_{r+1}}$ of this set we get a homogeneous linear equation

$$0 = f \oplus g(q_{s_1} \ldots q_{s_{r+1}}) = \\ g(q_{s_2} \cdots q_{s_{r+1}}) + g(q_{s_1} q_{s_3} \cdots q_{s_{r+1}}) + \cdots + g(q_{s_1} \cdots g_{s_r}) \tag{53}$$

The matrix of the homogeneous linear equation system formed by all these equations is the incidence matrix of r-subsets (of a set of N elements) into $r+1$-subsets. It has full rank [12]. Since it consists of $\binom{N}{r+1}$ equations and $\binom{N}{r}$ variables, we get that for sufficiently large N, the null-space is zero-dimensional, thus the homogeneous system has only the trivial solution. It follows, in particular, that (52) holds.

Thus, $g(m) = 0$ for all m, so f is a regular element.

Step 2. We construct infinitely many different regular non-units. Consider the element $\tilde{f}$, with

$$\tilde{f}(k) = \begin{cases} c_k & k \in \mathcal{PP} \\ 0 & \text{otherwise} \end{cases}$$

and where the c_k's are "sufficiently generic" non-zero complex numbers, then we claim that $\tilde{f}$, too, is a regular non-unit. With g, m, r as before, we have that, for $r = 0$,

$$0 = f \oplus g(p^a) == f(p^a)g(1) = c_{p^a} g(1).$$

We demand that $c_{p^a} \neq 0$, then $g(1) = 0$.

For a general r, we argue as follows: the incidence matrices that occurred before will be replaced with "generic" matrices whose elements are c_k's or zeroes, and which specialize, when setting all $c_k = 1$, to full-rank matrices. They must therefore have full rank, and the proof goes through.

Step 3. Let g be a unit in $\mathcal{A}$, and $\tilde{f}$ as above. We claim that if $g \oplus f$ is of the above form, i.e. supported on $\mathcal{PP}$, then g must be a constant. Hence there are infinitely many non-associate regular non-units of the above form.

To prove the claim, we argue exactly as before, using the fact that $g \oplus \tilde{f}$ is supported on $\mathcal{PP}$. For $m = q_1^{a_1} \cdots q_r^{a_r}$ as before, the case $r = 0$ yields nothing:

$$0 = g \oplus \tilde{f}(1) = \tilde{f}(1)g(1) = 0g(1) = 0,$$

neither does the case $r = 1$:

$$w = g \oplus \tilde{f}(q^a) = \tilde{f}(q^a)g(1),$$

so $g(1)$ may be non-zero. But for $r = 2$ we get

$$0 = g \oplus \tilde{f}(q_1^{a_1} q_2^{a_2}) = \tilde{f}(q_1^{a_1})g(q_2^{a_2}) + g(q_1^{a_1})\tilde{f}(q_2^{a_2}),$$

and also

$$0 = g \oplus \tilde{f}(q_1^{a_1} q_3^{a_3}) = \tilde{f}(q_1^{a_1})g(q_3^{a_3}) + g(q_1^{a_1})\tilde{f}(q_3^{a_3})$$
$$0 = g \oplus \tilde{f}(q_2^{a_2} q_3^{a_3}) = \tilde{f}(q_2^{a_2})g(q_3^{a_3}) + g(q_1^{a_1})\tilde{f}(q_3^{a_3})$$

which means that

$$\begin{bmatrix} \tilde{f}(q_2^{a_2}) & \tilde{f}(q_1^{a_1}) & 0 \\ \tilde{f}(q_3^{a_3}) & 0 & \tilde{f}(q_1^{a_1}) \\ 0 & \tilde{f}(q_3^{a_3}) & \tilde{f}(q_2^{a_2}) \end{bmatrix} \begin{bmatrix} g(q_1^{a_1}) \\ g(q_2^{a_2}) \\ g(q_3^{a_3}) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By our assumptions, the coefficient matrix is non-singular, so only the zero solution exists, hence $g(q_1^{a_1}) = 0$.

An analysis similar to what we did before shows that $g(q_1^{a_1} \cdots q_r^{a_r}) = 0$ for $r > 1$. $\square$

With the same method, one can easily show that the characteristic function on $\mathcal{P}$ is regular.

6. Some simple results on factorisation

Cashwell-Everett [4] showed that $(\mathcal{A}, +, \cdot)$ is a UFD. We will briefly treat the factorisation properties of $(\mathcal{A}, +, \oplus)$. Definitions and facts regarding factorisation in commutative rings with zero-divisors from the articles by Anderson and Valdes-Leon [1, 2] will be used.

First, we note that since $(\mathcal{A}, +, \oplus)$ is quasi-local, it is présimplifiable, i.e. $a \neq \mathbf{0}$, $a = r \oplus a$ implies that r is a unit. It follows that for $a, b \in \mathcal{A}$, the following three conditions are equivalent:

(1) a, b are *associates*, i.e. $\mathcal{A} \oplus a = \mathcal{A} \oplus b$.
(2) a, b are *strong associates*, i.e. $a = u \oplus b$ for some unit u.
(3) a, b are *very strong associates*, i.e. $\mathcal{A} \oplus a = \mathcal{A} \oplus b$ and either $a = b = \mathbf{0}$, or $a \neq \mathbf{0}$ and $a = r \oplus b \implies r \in U(\mathcal{A})$.

We say that $a \in \mathcal{A}$ is *irreducible*, or an *atom*, if $a = b \oplus c$ implies that a is associate with either b or c.

Theorem 6.1. $(\mathcal{A}, +, \oplus)$ *is* atomic*, i.e. all non-units can be written as a product of finitely many atoms. In fact,* $(\mathcal{A}, +, \oplus)$ *is a* bounded factorial ring *(BFR), i.e. there is a bound on the length of all factorisations of an element.*

Proof. It follows from Lemma 3.3 that the non-unit f has a factorisation into at most $\mathrm{D}(f)$ atoms. □

Example 6.2. We have that $e_2 \oplus (e_{2^k} + e_3) = e_6$ for all k, hence e_6 has an infinite number of non-associate irreducible divisors, and infinitely many factorisations into atoms.

Example 6.3. The element $h = e_{30}$ can be factored as $e_2 \oplus e_3 \oplus e_5$, or as $(e_6 + e_{20}) \oplus (e_2 + e_5)$.

These examples show that $(\mathcal{A}, +, \oplus)$ is neither a *half-factorial ring*, nor a *finite factorisation ring*, nor a *weak finite factorisation ring*, nor an *atomic idf-ring*.

7. The subring of arithmetical functions supported on square-free integers

Let $\mathcal{SQF} \subset \mathbb{N}^+$ denote the set of square-free integers, and put

$$\mathfrak{C} = \{\, f \in \mathcal{A} | \mathrm{supp}(f) \subset \mathcal{SQF} \,\} \tag{54}$$

For any $f \in \mathcal{A}$, denote by $p(f) \in \mathfrak{C}$ the restriction of f to $\mathcal{SQF}$.

Theorem 7.1. $(\mathfrak{C}, +, \oplus)$ *is a subring of* $(\mathcal{A}, +, \oplus)$*, and a closed* $\mathbb{C}$*-subalgebra with respect to the norm* $|\cdot|$*. The map*

$$\begin{aligned} p : \mathcal{A} &\to \mathfrak{C} \\ f &\mapsto p(f) \end{aligned} \tag{55}$$

is a continuous $\mathbb{C}$*-algebra epimorphism, and a retraction of the inclusion map* $\mathfrak{C} \subset \mathcal{A}$*.*

Proof. Let $f, g \in \mathfrak{C}$. If $n \in \mathbb{N}^+ \setminus \mathcal{SQF}$ then $(f+g)(n) = f(n) + g(n) = 0$, and $cf(n) = 0$ for all $c \in \mathbb{C}$. Since $n \in \mathbb{N}^+ \setminus \mathcal{SQF}$, there is at least on prime p such that $p^2 \,|n$. If m is a unitary divisor of m, then either m or n/m is divisible by p^2. Thus

$$(f \oplus g)(n) = \sum_{m||n} f(m)g(n/m) = 0.$$

If $f_k \to f$ in $\mathcal{A}$, and all $f_k \in \mathfrak{C}$, let $n \in \operatorname{supp}(f)$. Then there is an N such that $f(n) = f_k(n)$ for all $k \geq N$. But $\operatorname{supp}(f_k) \subset \mathcal{SQF}$, so $n \in \mathcal{SQF}$. This shows that $\mathfrak{C}$ is a closed subalgebra of $\mathcal{A}$.

It is clear that $p(f+g) = p(f) + p(g)$ and that $p(cf) = cp(f)$ for any $c \in \mathbb{C}$. If n is not square-free, we have already showed that

$$0 = (p(f) \oplus p(g))(n) = p((f \oplus g))(n).$$

Suppose therefore that n is square-free. Then so is all its unitary divisors, hence

$$p(f \oplus g)(n) = (f \oplus g)(n) = \sum_{m||n} f(m)g(n/m) = \sum_{m||n} p(f)(m)p(g)(n/m) = (p(f) \oplus p(g))(n).$$

We have that $p(f) = f$ if and only if $f \in \mathfrak{C}$, hence $p(p(f)) = p(f)$, so p is a retraction to the inclusion $i : \mathfrak{C} \to \mathcal{A}$. In other words, $p \circ i = \mathrm{id}_{\mathfrak{C}}$. □

Corollary 7.2. *The multiplicative inverse of an element in $\mathfrak{C}$ lies in $\mathfrak{C}$.*

Proof. If $f \in \mathfrak{C}$, $f \oplus g = e_1$ then

$$e_1 = p(e_1) = p(f \oplus g) = p(f) \oplus p(g) = f \oplus p(g),$$

hence $g = p(g)$, so $g \in \mathfrak{C}$.

Alternatively, we can reason as follows. If f is a unit in $\mathfrak{C}$ then we can without loss of generality assume that $f(1) = 1$. By Theorem 3.7, $g = -f + e_1$ is topologically nilpotent, hence by Proposition 1.2.4 of [3] we have that the inverse of $e_1 - g = f$ can be expressed as $\sum_{i=0}^{\infty} g^i$. It is clear that g, and every power of it, is supported on $\mathcal{SQF}$, hence so is f^{-1}. □

Corollary 7.3. $(\mathfrak{C}, +, \oplus)$ *is semi-local.*

Proof. The units consists of all $f \in \mathfrak{C}$ with $f(1) \neq 0$, and the non-units form the unique maximal ideal. □

Remark 7.4. More generally, given any subset $Q \subset \mathbb{N}^+$, we get a retract of $(\mathcal{A}, +, \oplus)$ when considering those arithmetical functions that are supported on the integers $n = p_1^{a_1} \cdots p_r^{a_r}$ with $a_i \in Q \cup \{0\}$. This property is unique for the unitary convolution, among all regular convolutions in the sense of Narkiewicz [7].

In particular, the set of arithmetical functions supported on the exponentially odd integers (those n for which all a_i are odd) forms a retract of $(\mathcal{A}, +, \oplus)$. It follows that the inverse of such a function is of the same form.

Let $T = \mathbb{C}[[x_1, x_2, x_3, \dots]]$, the large power series ring on countably many variables, and let J denote the ideal of elements supported on non square-free monomials.

Theorem 7.5. $(\mathfrak{C}, +, \oplus) \simeq T/J$. *This algebra can also be described as the generalized power series ring on the monoid-with-zero whose elements are all finite subsets of a fixed countable set X, with multiplication*

$$A \times B = \begin{cases} A \cup B & \text{if } A \cap B = \emptyset \\ 0 & \text{otherwise .} \end{cases} \tag{56}$$

Proof. Define η by

$$\eta : T \to \mathcal{A}$$
$$\eta(\sum_m c_m m) = \sum_{m \text{ square-free}} c_m e_m, \tag{57}$$

where for a square-free monomial $m = m_{i_1} \cdots m_{i_r}$ with $1 \le i_1 < \cdots < i_r$ we put $e_m = e_{p_{i_1} \cdots p_{i_r}}$. Then $\eta(T) = \mathfrak{C}$, $\ker \eta = J$. It follows that $\mathfrak{C} \simeq T/J$. $\square$

References

[1] D. D. Anderson and Silvia Valdes-Leon. Factorization in commutative rings with zero divisors. *Rocky Mountain J. Math.*, 26(2):439–480, 1996.

[2] D. D. Anderson and Silvia Valdes-Leon. Factorization in commutative rings with zero divisors. II. In *Factorization in integral domains (Iowa City, IA, 1996)*, pages 197–219. Dekker, New York, 1997.

[3] S. Bosch, U. Güntzer, and R. Remmert. *Non-Archimedean analysis*. Springer-Verlag, Berlin, 1984. A systematic approach to rigid analytic geometry.

[4] E. D. Cashwell and C. J. Everett. The ring of number-theorethic functions. *Pacific Journal of Mathematics*, 9:975–985, 1959.

[5] Eckford Cohen. Arithmetical functions associated with the unitary divisors of an integer. *Mathematische Zeitschrift*.

[6] James A. Huckaba. *Commutative rings with zero divisors*. Marcel Dekker Inc., New York, 1988.

[7] W. Narkiewicz. On a class of arithmetical convolutions. *Colloq. Math.*, 10:81–94, 1963.

[8] Emil D. Schwab and Gheorghe Silberberg. A note on some discrete valuation rings of arithmetical functions. *Archivum Mathematicum (Brno)*, 36:103–109, 2000.

[9] Emil D. Schwab and Gheorghe Silberberg. The valuated ring of the arithmetical functions as a power series ring. *Archivum Mathematicum (Brno)*, 37(1):77–80, 2001.

[10] R. Sivaramakrishnan. *Classical theory of arithmetic functions*, volume 126 of *Pure and applied mathematics*. Marcel Dekker, 1989.

[11] R. Vaidyanathaswamy. The theory of multiplicative arithmetic functions. *Trans. Amer. Math. Soc.*, 33(2):579–662, 1931.

[12] Richard M. Wilson. The necessary conditions for t-designs are sufficient for something. *Utilitas Mathematica*, 4:207–215, 1973.

[13] K. L. Yocom. Totally multiplicative functions in regular convolution rings. *Canad. Math. Bull.*, 16:119–128, 1973.

Department of Mathematics, Stockholm University, SE-10691 Stockholm, Sweden

E-mail address: `jans@matematik.su.se`